

Error analysis for the deep Kolmogorov method

Iulian Cîmpean^{1,2}, Thang Do³, Lukas Gonon⁴, Arnulf Jentzen^{5,6}, and Ionel Popescu^{7,8}

¹ Faculty of Mathematics and Computer Science,
University of Bucharest, Romania, e-mail: iulian.cimpean@unibuc.ro

² Institute of Mathematics “Simion Stoilow” of the Romanian Academy,
Romania, e-mail: iulian.cimpean@imar.ro

³ School of Data Science, The Chinese University of Hong Kong, Shenzhen
(CUHK-Shenzhen), China, e-mail: minhthangdo@link.cuhk.edu.cn

⁴ Faculty of Mathematics and Statistics, University of St. Gallen,
Switzerland, e-mail: lukas.gonon@unisg.ch

⁵ School of Data Science, The Chinese University of Hong Kong, Shenzhen
(CUHK-Shenzhen), China, e-mail: ajentzen@cuhk.edu.cn

⁶ Applied Mathematics: Institute for Analysis and Numerics, Faculty of Mathematics and
Computer Science, University of Münster, Germany, e-mail: ajentzen@uni-muenster.de

⁷ Faculty of Mathematics and Computer Science,
University of Bucharest, Romania, e-mail: ionel.popescu@fmi.unibuc.ro

⁸ Institute of Mathematics “Simion Stoilow” of the Romanian Academy,
Romania, e-mail: ionel.popescu@imar.ro

Abstract

The deep Kolmogorov method is a simple and popular deep learning based method for approximating solutions of partial differential equations (PDEs) of the Kolmogorov type. In this work we provide an error analysis for the deep Kolmogorov method for heat PDEs. Specifically, we reveal convergence with convergence rates for the overall mean square distance between the exact solution of the heat PDE and the realization function of the approximating deep neural network (DNN) associated with a stochastic optimization algorithm in terms of the size of the architecture (the depth/number of hidden layers and the width of the hidden layers) of the approximating DNN, in terms of the number of random sample points used in the loss function (the number of input-output data pairs used in the loss function), and in terms of the size of the optimization error made by the employed stochastic optimization method.

Contents

1	Introduction	2
1.1	Artificial neural networks (ANNs)	3
1.2	Error analysis for the deep Kolmogorov method	3
1.3	Literature review	5
1.4	Structure of this article	6
2	Explicit a priori estimates for ANNs	6
2.1	Explicit a priori estimates for realization functions of ANNs	6
2.2	Explicit a priori estimates for gradients of realization functions of ANNs	8

3	Error estimates for ANN approximations	10
3.1	Error estimates for shallow ANN approximations for C^2 -functions	10
3.2	Error estimates for deep ANN approximations for C^2 -functions	13
4	Error estimates for the deep Kolmogorov method	17
4.1	Sobolev embedding estimates with explicit constants	17
4.2	Feynman-Kac representations for solutions of heat PDEs	18
4.3	Error estimates for the Monte Carlo method	19
4.4	Error estimates for the deep Kolmogorov method	20
4.5	Convergence results for the deep Kolmogorov method	28

1. Introduction

It is nowadays a very active area of research to design and study deep learning methods to approximately solve *partial differential equations* (PDEs) and related scientific computing problems (cf., for example, the review articles and monographs [10, 27, 34, 41, 61, 67]). Popular such deep learning approximation methods for PDEs include *physics-informed neural networks* (PINNs) (cf. [90]) and the *deep Galerkin method* (DGM) (cf. [94]) as well as the deep *backward stochastic differential equation* (BSDE) method (cf. [33, 59]). Another quite simple and nowadays well-known deep learning approximation scheme is the *deep Kolmogorov method* (DKM) to approximately solve PDEs of the Kolmogorov type; see [9] and [23]. (In the references [9] and [23] the method has not been named as DKM but this has been done in the follow-up work [67, Chaper 17].) We also refer, for instance, to [8, 19] for suitable *deep splitting methods* (DSMs) that are based on the combination of the splitting method and the DKM.

Beyond the design and the numerical investigation of such deep learning approximation methods for PDEs, it is also an active topic of research to rigorously analyze such methods mathematically, that is, to establish rigorous upper bounds for the approximation error between the exact solution of the PDE and the deep learning approximation under consideration. While for classical approximation methods for solutions of PDEs such as finite difference (cf., for example, [72] and the references therein) and finite element methods (cf., for instance, [96] and the references therein) such error analyses for the difference between the PDE solution and the considered numerical approximation are – to a large extent – well understood, the situation is much more subtle in the case of deep learning approximation methods for PDEs and it basically remains a fundamental open problem of research to establish a full error analysis for any reasonable deep learning approximation scheme for PDEs.

There are, however, several partial error analyses for deep learning methods for PDEs in the literature. For instance, in the case of the PINNs Mishra & Molinaro [80, Theorem 3.1] explicitly bounds the overall root mean square distance between the exact solution of the PDE and the realization of the approximating *artificial neural network* (ANN) from above by the product of an error explicit constant and the sum of the loss function and the number of discretization points where the error constant also depends on the realization function of the approximating ANN. A qualitative convergence result without a convergence speed for the DKM applied to semilinear Kolmogorov PDEs in terms of the expectation of the loss function can also be found in [23, Proposition 4.7, Appendix A.3, and (21)].

In this work we push these error analyses, in the case of the DKM applied to heat PDEs, a bit further by establishing that there exists a strictly positive real error constant $c \in (0, \infty)$ that is completely independent of the number of random sample points in the loss function (the number of input-output data pairs in the loss function), completely independent of the architecture of the approximating deep ANNs (the depth/the number of hidden layers and the widths of the hidden layers of the approximation ANNs), and completely independent of the random ANN parameter vector computed with stochastic optimization methods such that the

overall mean square distance between the exact solution of the [PDE](#) and the random deep [ANN](#) approximation converges to zero with explicit convergence constants and rates using the error constant c as the *optimization error* of the underlying stochastic optimization algorithm converges to zero (convergence rate 1 in terms of the optimization error), as the number of *random* sample points used in the loss function (the number of input-output data pairs used in the loss function) converges to infinity (convergence rate $1/2$ in terms of the number of random sample points), and as the width of the approximating [ANN](#) (the number of neurons on the hidden layers of the approximating [ANN](#)) converges to infinity (convergence rate $2/(d+5)$ in terms of the width of the [ANN](#)); see (4) in [Theorem 1.2](#) below and [Corollary 4.9](#) in [Section 4](#) below. The specific analysis in this work is restricted to simple heat [PDEs](#), to [ANNs](#) with the *rectified linear unit* ([ReLU](#)) activation, and to the [DKM](#), which seems to be easier to analyze than many other deep learning approximation methods for [PDEs](#) such as [PINNs](#) and deep [BSDE](#) approximations, but we expect that our arguments can be extended to more challenging deep learning methods and [PDE](#) problems.

1.1. Artificial neural networks (ANNs)

To formulate our main error analysis result within this introductory section in [Theorem 1.2](#) below, we need to describe realization functions of [ANNs](#) in accurate mathematical terms. This is precisely the subject of the following notion in [Definition 1.1](#). In the simplest form, a realization function of an [ANN](#) can be considered as a function that is given by multiple compositions of affine functions and multidimensional versions of a certain fixed one-dimensional nonlinear function, which is referred to as activation function of the [ANN](#). In [Definition 1.1](#) the natural number $L \in \mathbb{N} = \{1, 2, 3, \dots\}$ specifies the number of affine functions in the compositions and the natural numbers $\ell_0, \ell_1, \dots, \ell_L$ specify the dimensionalities of the domains and codomains, respectively, of the affine functions in the compositions.

Definition 1.1 ([ANNs](#)). For every $L \in \mathbb{N}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in \mathbb{N}^{L+1}$ we denote by $\mathfrak{d}(\ell) \in \mathbb{N}$ the natural number given by $\mathfrak{d}(\ell) = \sum_{i=1}^L \ell_i(\ell_{i-1} + 1)$ and for every $L \in \mathbb{N}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in \mathbb{N}^{L+1}$, $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$ we denote by $\mathcal{N}_v^{\ell, \theta} = (\mathcal{N}_{v,1}^{\ell, \theta}, \dots, \mathcal{N}_{v,\ell_v}^{\ell, \theta}): \mathbb{R}^{\ell_0} \rightarrow \mathbb{R}^{\ell_v}$, $v \in \{1, 2, \dots, L\}$, the functions which satisfy for all $v \in \{0, 1, \dots, L-1\}$, $x = (x_1, \dots, x_{\ell_0}) \in \mathbb{R}^{\ell_0}$, $i \in \{1, 2, \dots, \ell_{v+1}\}$ that

$$\begin{aligned} \mathcal{N}_{v+1,i}^{\ell, \theta}(x) &= \theta_{\ell_{v+1}\ell_v + i + \sum_{h=1}^v \ell_h(\ell_{h-1} + 1)} \\ &+ \sum_{j=1}^{\ell_v} \theta_{(i-1)\ell_v + j + \sum_{h=1}^v \ell_h(\ell_{h-1} + 1)} (x_j \mathbb{1}_{\{0\}}(v) + \max\{\mathcal{N}_{\max\{v,1\},j}^{\ell, \theta}(x), 0\} \mathbb{1}_{\mathbb{N}}(v)). \end{aligned} \quad (1)$$

In (1) appears the [ReLU](#) activation function $\mathbb{R} \ni x \mapsto \max\{x, 0\} \in \mathbb{R}$ and in [Definition 1.1](#) we thus describe [ANNs](#) with the [ReLU](#) activation function. [Definition 1.1](#) essentially coincides with the formulation of [ANNs](#) in, for example, [[30](#), Setting 2.1].

1.2. Error analysis for the deep Kolmogorov method

Having specified realization functions of [ReLU ANNs](#) in [Definition 1.1](#) above, we are now in the position to formulate in [Theorem 1.2](#) below one of the main error analysis results of this work. In [Theorem 1.2](#) the function $u: [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ is the solution of the heat equation in (2) that we intend to approximate by deep [ANNs](#) by means of the [DKM](#); see (3) for the loss function of the [DKM](#) in which *independent and identically distributed* (i.i.d.) random variables appear. We now present [Theorem 1.2](#) with all mathematical details and, thereafter, we briefly explain the contribution of [Theorem 1.2](#) and some of the mathematical objects appearing in [Theorem 1.2](#) in words.

Theorem 1.2. Let $T \in (0, \infty)$, $d \in \mathbb{N}$, let $u \in C^2([0, T] \times \mathbb{R}^d, \mathbb{R})$ be at most polynomially growing¹, assume for all $t \in (0, T)$, $x \in \mathbb{R}^d$ that

$$\frac{\partial}{\partial t} u(t, x) + \frac{1}{2} \Delta_x u(t, x) = 0, \quad (2)$$

let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $\mathbb{T}_n: \Omega \rightarrow [0, T]$, $n \in \mathbb{N}$, be independent uniformly random variables, let $D \subseteq \mathbb{R}^d$ be open and bounded, let $\mathbb{X}_n: \Omega \rightarrow D$, $n \in \mathbb{N}$, be independent uniformly distributed random variables, let $W^n: [0, T] \times \Omega \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}$, be *i.i.d.* standard Brownian motions, assume that $(\mathbb{T}_n)_{n \in \mathbb{N}}$, $(\mathbb{X}_n)_{n \in \mathbb{N}}$, and $(W^n)_{n \in \mathbb{N}}$ are independent, and for every $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ let $\mathbb{F}_{M, \ell}: \mathbb{R}^{\mathfrak{d}(\ell)} \rightarrow \mathbb{R}$ satisfy for all $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ that

$$\mathbb{F}_{M, \ell}(\theta) = \frac{1}{M_1} \sum_{m=1}^{M_1} |\mathcal{N}_L^{\ell, \theta}(\mathbb{T}_m, \mathbb{X}_m) - [\frac{1}{M_2} \sum_{n=1}^{M_2} u(T, \mathbb{X}_m + W_{T-\mathbb{T}_m}^{m M_2 + n})]|^2. \quad (3)$$

Then there exists $c \in (0, \infty)$ such that for every $R \in [c, \infty)$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ and every random variable $\vartheta: \Omega \rightarrow [-R, R]^{\mathfrak{d}(\ell)}$ it holds that

$$\begin{aligned} \mathbb{E}[\int_{[0, T] \times D} |u(y) - \mathcal{N}_L^{\ell, \vartheta}(y)|^2 dy] &\leq [R + \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{cL} (M_1)^{-1/2} \\ &+ c[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{-2/(d+5)} + c \mathbb{E}[\mathbb{F}_{M, \ell}(\vartheta) - \inf_{\theta \in [-R, R]^{\mathfrak{d}(\ell)}} \mathbb{F}_{M, \ell}(\theta)] \end{aligned} \quad (4)$$

(cf. Definition 1.1).

Theorem 1.2 is an immediate consequence of Corollary 4.8 in Subsection 4.4 below. Corollary 4.8, in turn, is established through an application of Theorem 4.6 in Section 4 below, which is the main result of this work (see Subsection 4.4 for details).

The strictly positive real number $T \in (0, \infty)$ in Theorem 1.2 describes the time horizon of the heat equation under consideration and the natural number $d \in \mathbb{N}$ specifies the space dimension of the heat equation under consideration. In (2) we present the heat PDE whose solutions we intend to approximate by means of the DKM in (3) and the function $u: [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a solution of the PDE in (2). The function specified in (3) is precisely the loss function of the DKM (cf., for instance, [9, Framework 3.2] and [23, Subsection 4.2]). The vector $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ in (3) and (4) specifies the architecture of the considered approximating ANNs in the sense

- that ℓ_0 specifies the number of neurons on the 1st layer of the ANN (the number of neurons on the input layer),
- that ℓ_1 specifies the number of neurons on the 2nd layer of the ANN (the number of neurons on the 1st hidden layer),
- that ℓ_2 specifies the number of neurons on the 3rd layer of the ANN (the number of neurons on the 2nd hidden layer),
- ... ,
- that ℓ_{L-1} specifies the number of neurons on the L^{th} layer of the ANN (the number of neurons on the $(L-1)^{\text{th}}$ hidden layer), and
- that ℓ_L specifies the number of neurons on the $(L+1)^{\text{th}}$ layer of the ANN (the number of neurons on the output layer).

¹Note that for every $d \in \mathbb{N}$, every $A \subseteq \mathbb{R}^d$, and every $f: A \rightarrow \mathbb{R}$ it holds that f is at most polynomially growing if and only if there exists $c \in \mathbb{R}$ such that for all $x = (x_1, \dots, x_d) \in A$ it holds that $|f(x)| \leq c(1 + \sum_{i=1}^d |x_i|)^c$.

We note that (4) in Theorem 1.2 assures that there exists a strictly positive real error constant $c \in (0, \infty)$ such that for every upper bound $R \in [c, \infty)$ for the parameters of the ANN associated to the stochastic optimization method, every number of random sample points $M_1 \in \mathbb{N}$ in the loss function (see (3)), every number of Monte Carlo samples $M_2 \in \mathbb{N}$ in the loss function (see (3)), every depth $L \in \mathbb{N} \setminus \{1\}$ of the approximating ANNs, every architecture vector $\ell = (\ell_0, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ for the approximating ANNs, and every random variable $\vartheta: \Omega \rightarrow [-R, R]^{\mathfrak{d}(\ell)}$ resulting from the stochastic optimization method we have that the overall mean squared distance

$$\mathbb{E}[\int_D |u(y) - \mathcal{N}_L^{\ell, \vartheta}(y)|^2 dy] \tag{5}$$

on the bounded set D between the exact solution $u: [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ of the PDE in (2) and the realization function $\mathcal{N}_L^{\ell, \vartheta}: \mathbb{R}^{d+1} \rightarrow \mathbb{R}$ of the ANN associated with the random ANN parameter vector ϑ is bounded by the sum of the term $[R + \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{cL} (M_1)^{-1/2}$ (that converges to zero as M_1 increases to infinity), of the term $c[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{-2/(d+5)}$ (that converges to zero as the minimum $\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}$ increases to infinity), and of – up to the constant c – the optimization error $\mathbb{E}[\mathbb{F}_{M, \ell}(\vartheta) - \inf_{\theta \in [-R, R]^{\mathfrak{d}(\ell)}} \mathbb{F}_{M, \ell}(\theta)]$.

1.3. Literature review

In this subsection we provide a short review on selected further results in the literature that develop error analyses for deep learning based approximation schemes for PDEs and related problems. For more comprehensive reviews on deep learning approximation methods for PDEs in the literature we refer, for instance, to the overview and review articles [10, 17, 20, 27, 34, 41, 48, 61] and the references therein.

The mathematical analysis of deep learning methods for solving PDEs has attracted increasing attention in recent years. Numerous works have contributed insights into the approximation capabilities and theoretical guarantees of such methods. Substantial progress has been made on bounding individual components of the total error – such as approximation, sampling, or optimization errors in the past years. Several works in the literature focus on analysing the approximation errors for neural network approximations to PDE solutions for heat equations [47], Black Scholes PDEs [37, 52], non-local PDEs [7, 49, 50, 97], parametric PDEs [75], smoothness spaces associated to PDEs [1, 12, 51, 56, 86], semilinear (heat) equations [2, 3, 64, 65, 82], non-linear waves [81], zero-sum games [91], stochastic volatility models [15], optimal stopping problems [46], or McKean-Vlasov equations [83]. Several articles analyse approximation and sampling errors jointly by focusing on the empirical risk minimization problem (see, for example, [13, 14, 102]). Concerning the analysis of the generalization error we refer to [78] for the analysis of the generalization error for deep learning-based approximations for certain type of second-order linear PDEs and two-layer networks. For an analysis of the generalization error of several related deep learning methods for elliptic PDEs we refer to [53] for deep mixed residual methods, to [35, 69, 76] for the deep Ritz method, and to [70] for a DGM for weak solutions. Full error analysis for random feature neural network-based methods are provided in [45, 84, 85]. An error analysis for an ANN-based algorithm for ergodic mean field control is provided in [21].

In [78, Theorem 2.4] and [80, Theorem 3.1] Mishra and Molinaro provide upper estimates for the generalization/overall error of PINNs for approximating PDEs and inverse problems with error constants depending on the approximating (ANN realization) function. The error bounds in [28] provide separate analyses of the relevant error components. [25] propose and mathematically analyse Friedrichs learning as a novel deep learning methodology that can learn the weak solutions of PDEs via a minimax formulation.

Moreover, there are various convergence results for minimizers of loss functions of deep learning algorithms in the literature focusing on discretization or simulation errors, without explicit approximation rates for the underlying ANNs or bounds in terms of the optimization error. For example, the following articles provide such convergence results for the deep BSDE method and

related (forward) BSDE based methods [5, 6, 16, 24, 60, 68, 73, 88, 98, 100, 104], deep backward schemes [22, 42, 63, 87], the deep splitting scheme [39, 85], the deep fictitious play algorithm [58], algorithms for first order *Hamilton-Jacobi-Bellman* (HJB) equations [18], methods related to the DGM [43, 94], PINNs [4, 40, 62, 71], and a randomized quasi-Monte Carlo based deep learning method [101].

Results for related algorithms are provided, for instance, for regularized dynamical parametric approximations in [38] for the variational Monte Carlo algorithm in [36], for physics-informed kernel learning in [31, 32], for a Monte Carlo Euler scheme in [26], for a quasi-Regression Monte Carlo Scheme in [44], and for a policy iteration algorithm [99].

Finally, we also refer, for example, to [92, Lemma 4] and [93, Appendix C] for abstract general error estimates for ANN approximations in terms of the optimization error.

1.4. Structure of this article

The remainder of this article is structured in the following way. After this introductory section, we establish in Section 2 several elementary explicit a priori estimates for realization functions for ANNs. In Section 3 we present in Corollary 3.7 suitable approximation error estimates for deep ANN approximations of twice continuously differentiable functions. Finally, in Section 4 we combine the findings from Section 2 and Section 3 to establish in Proposition 4.5, Theorem 4.6, Corollary 4.7, and Corollary 4.8 error estimates for the deep Kolmogorov method. Theorem 1.2 above is a direct consequence of Corollary 4.8.

2. Explicit a priori estimates for ANNs

In this section we establish in Lemma 2.2, Lemma 2.3, and Corollary 2.4 several elementary a priori estimates for realizations of arbitrarily deep ANNs with general activation functions. The arguments in our proofs of Lemma 2.2, Lemma 2.3, and Corollary 2.4 are based, for instance, on [11, Subsection 2.1.5] and [14, Theorem 2.6 and Section A.3], respectively. In Section 4 we employ the a priori estimates in Lemma 2.2 and Corollary 2.4 to establish in Proposition 4.5 error estimates for the deep Kolmogorov method and, thereby, to prove Theorem 1.2 in the introduction.

2.1. Explicit a priori estimates for realization functions of ANNs

Setting 2.1. Let $d, L \in \mathbb{N}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d\} \times \mathbb{N}^{L-1} \times \{1\})$, for every $k \in \{0, 1, \dots, L\}$ let $\mathbf{d}_k \in \mathbb{R}$ satisfy $\mathbf{d}_k = \sum_{h=1}^k \ell_h(\ell_{h-1} + 1)$, let $\mathbb{A}: \mathbb{R} \rightarrow \mathbb{R}$ be a function, and for every $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$ let $\mathbf{N}_v^\theta = (\mathbf{N}_{v,1}^\theta, \dots, \mathbf{N}_{v,\ell_v}^\theta): \mathbb{R}^{\ell_0} \rightarrow \mathbb{R}^{\ell_v}$, $v \in \{1, 2, \dots, L\}$, satisfy for all $v \in \{0, 1, \dots, L-1\}$, $x = (x_1, \dots, x_{\ell_0}) \in \mathbb{R}^{\ell_0}$, $i \in \{1, 2, \dots, \ell_{v+1}\}$ that

$$\begin{aligned} \mathbf{N}_{v+1,i}^\theta(x) &= \theta_{\ell_{v+1}\ell_v+i+\mathbf{d}_v} \\ &\quad + \sum_{j=1}^{\ell_v} \theta_{(i-1)\ell_v+j+\mathbf{d}_v} (x_j \mathbb{1}_{\{0\}}(v) + \mathbb{A}(\mathbf{N}_{\max\{v,1\},j}^\theta(x)) \mathbb{1}_{\mathbb{N}}(v)) \end{aligned} \quad (6)$$

(cf. Definition 1.1).

Lemma 2.2. Assume Setting 2.1 and let $q \in [0, \infty)$, $c, a \in \mathbb{R}$, $b \in (a, \infty)$ satisfy for all $x \in \mathbb{R}$ that $|\mathbb{A}(x)| \leq c(1 + |x|^q)$. Then it holds for all $k \in \{1, 2, \dots, L\}$, $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$ that

$$\begin{aligned} &\max_{i \in \{1, 2, \dots, \ell_k\}} \sup_{x \in [a, b]^d} |\mathbf{N}_{k,i}^\theta(x)| \\ &\leq [\max\{1, |a|, |b|\}]^{\max\{1, q^{k-1}\}} [\max\{1, \max_{j \in \{1, 2, \dots, \mathbf{d}_k\}} |\theta_j|\}]^{k \max\{1, q^{k-1}\}} \\ &\quad \cdot [1 + 2 \max\{c, 1\} \max\{\ell_0, \ell_1, \dots, \ell_{k-1}\}]^{k \max\{1, q^{k-1}\}}. \end{aligned} \quad (7)$$

Proof of Lemma 2.2. We prove (7) by induction on $k \in \{1, 2, \dots, L\}$. For the base case $k = 1$ note that (6) shows that for all $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $x = (x_1, \dots, x_d) \in [a, b]^d$, $i \in \{1, 2, \dots, \ell_1\}$ it holds that

$$\begin{aligned} |\mathbf{N}_{1,i}^\theta(x)| &= |\theta_{\ell_1 \ell_0 + i} + \sum_{j=0}^d \theta_{(i-1)\ell_0 + j} x_j| \leq |\theta_{\ell_1 \ell_0 + i}| + \left| \sum_{j=0}^d \theta_{(i-1)\ell_0 + j} x_j \right| \\ &\leq (d+1) [\max\{1, |a|, |b|\} \max_{j \in \{1, 2, \dots, \mathbf{d}_1\}} |\theta_j|] \\ &\leq [\max\{1, |a|, |b|\}]^{\max\{1, q^0\}} [\max\{1, \max_{j \in \{1, 2, \dots, \mathbf{d}_{0+1}\}} |\theta_j|\}]^{(0+1) \max\{1, q^0\}} \\ &\quad \cdot [1 + 2 \max\{c, 1\} \ell_0]^{(0+1) \max\{1, q^k\}}. \end{aligned} \tag{8}$$

This proves (7) in the base case $k = 1$. For the induction step we assume that there exists $k \in \mathbb{N} \cap [0, L)$ which satisfies for all $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $x \in [a, b]^d$ that

$$\begin{aligned} \max_{i \in \{1, 2, \dots, \ell_k\}} |\mathbf{N}_{k,i}^\theta(x)| &\leq [\max\{1, |a|, |b|\}]^{\max\{1, q^{k-1}\}} [\max\{1, \max_{j \in \{1, 2, \dots, \mathbf{d}_k\}} |\theta_j|\}]^{k \max\{1, q^{k-1}\}} \\ &\quad \cdot [1 + 2 \max\{c, 1\} \max\{\ell_0, \ell_1, \dots, \ell_k\}]^{k \max\{1, q^{k-1}\}}. \end{aligned} \tag{9}$$

Observe that (6) ensures that for all $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $x \in [a, b]^d$, $i \in \{1, 2, \dots, \ell_{k+1}\}$ it holds that

$$\begin{aligned} |\mathbf{N}_{k+1,i}^\theta(x)| &\leq |\theta_{\ell_{k+1} \ell_k + i + \mathbf{d}_k}| + \left[\sum_{j=1}^{\ell_k} |\theta_{(i-1)\ell_k + j + \mathbf{d}_k} \mathbb{A}(\mathbf{N}_{k,j}^\theta(x))| \right] \\ &\leq [\max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j|] \\ &\quad + \ell_k [\max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j|] [\max_{j \in \{1, 2, \dots, \ell_k\}} |\mathbb{A}(\mathbf{N}_{k,j}^\theta(x))|]. \end{aligned} \tag{10}$$

Combining this and the assumption that for all $x \in \mathbb{R}$ it holds that $|\mathbb{A}(x)| \leq c(1 + |x|^q)$ implies that for all $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $x \in [a, b]^d$, $i \in \{1, 2, \dots, \ell_{k+1}\}$ it holds that

$$\begin{aligned} |\mathbf{N}_{k+1,i}^\theta(x)| &\leq [\max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j|] \\ &\quad + \ell_k [\max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j|] [\max_{j \in \{1, 2, \dots, \ell_k\}} (c + c |\mathbf{N}_{k,j}^\theta(x)|^q)] \\ &= [\max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j|] (1 + \ell_k c + \ell_k c [\max_{j \in \{1, 2, \dots, \ell_k\}} |\mathbf{N}_{k,j}^\theta(x)|^q]). \end{aligned} \tag{11}$$

This and (9) show for all $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $x \in [a, b]^d$, $i \in \{1, 2, \dots, \ell_{k+1}\}$ that

$$\begin{aligned} |\mathbf{N}_{k+1,i}^\theta(x)| &\leq (1 + \ell_k c) [\max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j|] \\ &\quad + \ell_k c [\max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j|] [\max\{1, |a|, |b|\}]^{q \max\{1, q^{k-1}\}} \\ &\quad \cdot [\max\{1, \max_{j \in \{1, 2, \dots, \mathbf{d}_k\}} |\theta_j|\}]^{qk \max\{1, q^{k-1}\}} \\ &\quad \cdot [1 + 2 \max\{c, 1\} \max\{\ell_1, \ell_2, \dots, \ell_{k-1}\}]^{qk \max\{1, q^{k-1}\}} \\ &\leq (1 + \ell_k c) [\max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j|] \\ &\quad + \ell_k c [\max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j|] [\max\{1, |a|, |b|\}]^{\max\{1, q^k\}} \\ &\quad \cdot [\max\{1, \max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j|\}]^{k \max\{1, q^k\}} \\ &\quad \cdot [1 + 2 \max\{c, 1\} \max\{\ell_1, \ell_2, \dots, \ell_k\}]^{k \max\{1, q^k\}}. \end{aligned} \tag{12}$$

Hence, we obtain for all $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $x \in [a, b]^d$, $i \in \{1, 2, \dots, \ell_{k+1}\}$ that

$$\begin{aligned} |\mathbf{N}_{k+1,j}^\theta(x)| &\leq (1 + 2\ell_k c) [\max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j|] [\max\{1, |a|, |b|\}]^{\max\{1, q^k\}} \\ &\quad \cdot [\max\{1, \max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j|\}]^{k \max\{1, q^k\}} \\ &\quad \cdot [1 + 2 \max\{c, 1\} \max\{\ell_1, \ell_2, \dots, \ell_k\}]^{k \max\{1, q^k\}} \\ &\leq [\max\{1, |a|, |b|\}]^{\max\{1, q^k\}} [\max\{1, \max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j|\}]^{(k+1) \max\{1, q^k\}} \\ &\quad \cdot [1 + 2 \max\{c, 1\} \max\{\ell_1, \ell_2, \dots, \ell_k\}]^{(k+1) \max\{1, q^k\}}. \end{aligned} \tag{13}$$

Combining (8) and induction therefore proves (7). The proof of Lemma 2.2 is thus complete. \square

2.2. Explicit a priori estimates for gradients of realization functions of ANNs

Lemma 2.3. *Assume Setting 2.1 and let $c, a \in \mathbb{R}$, $b \in (a, \infty)$ satisfy for all $x, y \in \mathbb{R}$ that $|\mathbb{A}(x) - \mathbb{A}(y)| \leq c|x - y|$. Then it holds for all $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)})$, $\vartheta = (\vartheta_1, \dots, \vartheta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $x \in [a, b]^d$ that*

$$\begin{aligned} |\mathbf{N}_L^\theta(x) - \mathbf{N}_L^\vartheta(x)| &\leq \max\{1, |a|, |b|\} [(3 + |\mathbb{A}(0)|) \max\{c, 1\} \max\{\ell_0, \ell_1, \dots, \ell_{L-1}\}]^L \\ &\quad \cdot [\max_{j \in \{1, 2, \dots, \mathfrak{d}(\ell)\}} \max\{1, |\theta_j|, |\vartheta_j|\}]^L [\max_{j \in \{1, 2, \dots, \mathfrak{d}(\ell)\}} |\theta_j - \vartheta_j|]. \end{aligned} \quad (14)$$

Proof of Lemma 2.3. Throughout this proof for every $k \in \{0, 1, \dots, L\}$ let $\mathbf{d}_k \in \mathbb{R}$ satisfy $\mathbf{d}_k = \sum_{m=1}^k \ell_m(\ell_{m-1} + 1)$. We claim that for all $k \in \{1, 2, \dots, L\}$, $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)})$, $\vartheta = (\vartheta_1, \dots, \vartheta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $x \in [a, b]^d$ it holds that

$$\begin{aligned} &\max_{i \in \{1, 2, \dots, \ell_k\}} |\mathbf{N}_{k,i}^\theta(x) - \mathbf{N}_{k,i}^\vartheta(x)| \\ &\leq \max\{1, |a|, |b|\} [\max_{j \in \{1, 2, \dots, \mathbf{d}_k\}} \max\{1, |\theta_j|, |\vartheta_j|\}]^k \\ &\quad \cdot [1 + (2 + |\mathbb{A}(0)|) \max\{c, 1\} \max\{\ell_0, \ell_1, \dots, \ell_{k-1}\}]^k [\max_{j \in \{1, 2, \dots, \mathbf{d}_k\}} |\theta_j - \vartheta_j|]. \end{aligned} \quad (15)$$

We prove (15) by induction on $k \in \{1, 2, \dots, L\}$. For the base case $k = 1$ note that (1) shows that for all $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)})$, $\vartheta = (\vartheta_1, \dots, \vartheta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $x = (x_1, \dots, x_d) \in [a, b]^d$, $i \in \{1, 2, \dots, \ell_1\}$ it holds that

$$\begin{aligned} |\mathbf{N}_{1,i}^\theta(x) - \mathbf{N}_{1,i}^\vartheta(x)| &= |\theta_{\ell_1 \ell_0 + i} + \sum_{j=0}^d \theta_{(i-1)\ell_0 + j} x_j - (\vartheta_{\ell_1 \ell_0 + i} + \sum_{j=0}^d \vartheta_{(i-1)\ell_0 + j} x_j)| \\ &\leq |\theta_{\ell_1 \ell_0 + i} - \vartheta_{\ell_1 \ell_0 + i}| + |\sum_{j=0}^d (\theta_{(i-1)\ell_0 + j} - \vartheta_{(i-1)\ell_0 + j}) x_j| \\ &\leq (d + 1) [\max\{1, |a|, |b|\}] [\max_{j \in \{1, 2, \dots, \mathbf{d}_1\}} |\theta_j - \vartheta_j|] \\ &\leq [\max\{1, |a|, |b|\}] [\max_{j \in \{1, 2, \dots, \mathbf{d}_{0+1}\}} \max\{1, |\theta_j|, |\vartheta_j|\}]^1 \\ &\quad \cdot [1 + (2 + |\mathbb{A}(0)|) \max\{c, 1\} \ell_0]^1 [\max_{j \in \{1, 2, \dots, \mathbf{d}_1\}} |\theta_j - \vartheta_j|]. \end{aligned} \quad (16)$$

This proves (15) in the base case $k = 1$. For the induction step we assume that there exists $k \in \mathbb{N} \cap [0, L)$ which satisfies for all $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)})$, $\vartheta = (\vartheta_1, \dots, \vartheta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $x \in [a, b]^d$ that

$$\begin{aligned} &\max_{i \in \{1, 2, \dots, \ell_k\}} |\mathbf{N}_{k,i}^\theta(x) - \mathbf{N}_{k,i}^\vartheta(x)| \\ &\leq \max\{1, |a|, |b|\} [\max_{j \in \{1, 2, \dots, \mathbf{d}_k\}} \max\{1, |\theta_j|, |\vartheta_j|\}]^k \\ &\quad \cdot [1 + (2 + |\mathbb{A}(0)|) \max\{c, 1\} \max\{\ell_0, \ell_1, \dots, \ell_{k-1}\}]^k [\max_{j \in \{1, 2, \dots, \mathbf{d}_k\}} |\theta_j - \vartheta_j|]. \end{aligned} \quad (17)$$

Note that (6) demonstrates that for all $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)})$, $\vartheta = (\vartheta_1, \dots, \vartheta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $x \in [a, b]^d$, $i \in \{1, 2, \dots, \ell_{k+1}\}$ it holds that

$$\begin{aligned} &|\mathbf{N}_{k+1,i}^\theta(x) - \mathbf{N}_{k+1,i}^\vartheta(x)| \\ &\leq |\theta_{\ell_{k+1} \ell_k + i + \mathbf{d}_k} - \vartheta_{\ell_{k+1} \ell_k + i + \mathbf{d}_k}| \\ &\quad + [\sum_{j=1}^{\ell_k} |\theta_{(i-1)\ell_k + j + \mathbf{d}_k} \mathbb{A}(\mathbf{N}_{k,j}^\theta(x)) - \vartheta_{(i-1)\ell_k + j + \mathbf{d}_k} \mathbb{A}(\mathbf{N}_{k,j}^\vartheta(x))|] \\ &\leq [\max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j - \vartheta_j|] \\ &\quad + \ell_k [\max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j|] [\max_{j \in \{1, 2, \dots, \ell_k\}} |\mathbb{A}(\mathbf{N}_{k,j}^\theta(x)) - \mathbb{A}(\mathbf{N}_{k,j}^\vartheta(x))|] \\ &\quad + \ell_k [\max_{j \in \{1, 2, \dots, \mathbf{d}_{k+1}\}} |\theta_j - \vartheta_j|] [\max_{j \in \{1, 2, \dots, \ell_k\}} |\mathbb{A}(\mathbf{N}_{k,j}^\vartheta(x))|]. \end{aligned} \quad (18)$$

The assumption that for all $x, y \in \mathbb{R}$ it holds that $|\mathbb{A}(x) - \mathbb{A}(y)| \leq c|x - y|$ hence establishes that for all $i \in \{1, 2, \dots, \ell_{k+1}\}$, $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)})$, $\vartheta = (\vartheta_1, \dots, \vartheta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $x \in [a, b]^d$ it holds that

$$\begin{aligned}
& |\mathbf{N}_{k+1,i}^\theta(x) - \mathbf{N}_{k+1,i}^\vartheta(x)| \\
& \leq [\max_{j \in \{1, 2, \dots, \mathfrak{d}_{k+1}\}} |\theta_j - \vartheta_j|] \\
& + \ell_k c [\max_{j \in \{1, 2, \dots, \mathfrak{d}_{k+1}\}} |\theta_j|] [\max_{j \in \{1, 2, \dots, \ell_k\}} |\mathbf{N}_{k,j}^\theta(x) - \mathbf{N}_{k,j}^\vartheta(x)|] \\
& + \ell_k |\mathbb{A}(0)| [\max_{j \in \{1, 2, \dots, \mathfrak{d}_{k+1}\}} |\theta_j - \vartheta_j|] \\
& + \ell_k c [\max_{j \in \{1, 2, \dots, \mathfrak{d}_{k+1}\}} |\theta_j - \vartheta_j|] [\max_{j \in \{1, 2, \dots, \ell_k\}} |\mathbf{N}_{k,j}^\vartheta(x)|].
\end{aligned} \tag{19}$$

This, (17), and Lemma 2.2 ensure for all $i \in \{1, 2, \dots, \ell_{k+1}\}$, $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)})$, $\vartheta = (\vartheta_1, \dots, \vartheta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $x \in [a, b]^d$ that

$$\begin{aligned}
& |\mathbf{N}_{k+1,i}^\theta(x) - \mathbf{N}_{k+1,i}^\vartheta(x)| \\
& \leq (1 + \ell_k |\mathbb{A}(0)|) [\max_{j \in \{1, 2, \dots, \mathfrak{d}_{k+1}\}} |\theta_j - \vartheta_j|] \\
& + \ell_k c [\max_{j \in \{1, 2, \dots, \mathfrak{d}_{k+1}\}} |\theta_j|] \max\{1, |a|, |b|\} [\max_{j \in \{1, 2, \dots, \mathfrak{d}_k\}} \max\{1, |\theta_j|, |\vartheta_j|\}]^k \\
& \cdot [\max_{l \in \{0, 1, \dots, k-1\}} (1 + (2 + |\mathbb{A}(0)|) \max\{c, 1\} \ell_l)]^k [\max_{j \in \{1, 2, \dots, \mathfrak{d}_k\}} |\theta_j - \vartheta_j|] \\
& + \ell_k c [\max_{j \in \{1, 2, \dots, \mathfrak{d}_{k+1}\}} |\theta_j - \vartheta_j|] [\max\{1, |a|, |b|\}] \\
& \cdot [\max\{1, \max_{j \in \{1, 2, \dots, \mathfrak{d}(\ell)\}} |\vartheta_j|\}]^k [1 + 2 \max\{c, 1\} \max\{\ell_0, \ell_1, \dots, \ell_{L-1}\}]^k.
\end{aligned} \tag{20}$$

Hence, we obtain for all $i \in \{1, 2, \dots, \ell_{k+1}\}$, $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)})$, $\vartheta = (\vartheta_1, \dots, \vartheta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $x \in [a, b]^d$ that

$$\begin{aligned}
& |\mathbf{N}_{k+1,i}^\theta(x) - \mathbf{N}_{k+1,i}^\vartheta(x)| \\
& \leq [\max_{j \in \{1, 2, \dots, \mathfrak{d}_{k+1}\}} |\theta_j - \vartheta_j|] [\max\{1, |a|, |b|\}] [\max_{j \in \{1, 2, \dots, \mathfrak{d}_k\}} \max\{1, |\theta_j|, |\vartheta_j|\}]^k \\
& \cdot [1 + (2 + |\mathbb{A}(0)|) \max\{c, 1\} \max\{\ell_0, \ell_1, \dots, \ell_{k-1}\}]^k \\
& \cdot [1 + \ell_k |\mathbb{A}(0)| + \ell_k c [\max_{j \in \{1, 2, \dots, \mathfrak{d}_{k+1}\}} |\theta_j|] + \ell_k c] \\
& \leq [\max_{j \in \{1, 2, \dots, \mathfrak{d}_{k+1}\}} |\theta_j - \vartheta_j|] [\max\{1, |a|, |b|\}] [\max_{j \in \{1, 2, \dots, \mathfrak{d}_k\}} \max\{1, |\theta_j|, |\vartheta_j|\}]^k \\
& \cdot [1 + (2 + |\mathbb{A}(0)|) \max\{c, 1\} \max\{\ell_0, \ell_1, \dots, \ell_{k-1}\}]^k \\
& \cdot [1 + \ell_k |\mathbb{A}(0)| + 2\ell_k c] [\max\{1, \max_{j \in \{1, 2, \dots, \mathfrak{d}_{k+1}\}} |\theta_j|\}] \\
& \leq [\max_{j \in \{1, 2, \dots, \mathfrak{d}_{k+1}\}} |\theta_j - \vartheta_j|] [\max\{1, |a|, |b|\}] [\max_{j \in \{1, 2, \dots, \mathfrak{d}_k\}} \max\{1, |\theta_j|, |\vartheta_j|\}]^{k+1} \\
& \cdot [1 + (2 + |\mathbb{A}(0)|) \max\{c, 1\} \max\{\ell_0, \ell_1, \dots, \ell_k\}]^{k+1}.
\end{aligned} \tag{21}$$

Combining (16) and induction therefore implies (15). Observe that (15) shows (14). The proof of Lemma 2.3 is thus complete. \square

Corollary 2.4. *Assume Setting 2.1, let $a \in \mathbb{R}$, $b \in (a, \infty)$, and assume $\mathbb{A} \in C^1(\mathbb{R}, \mathbb{R})$. Then it holds² for all $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $x \in [a, b]^d$ that*

$$\begin{aligned}
& \|\nabla_\theta(\mathbf{N}_L^\theta(x))\| \leq \max\{1, |a|, |b|\} (3 + |\mathbb{A}(0)|)^L \\
& \cdot [\max\{\ell_0, \ell_1, \dots, \ell_{L-1}\} \max\{1, \max_{j \in \{1, 2, \dots, \mathfrak{d}(\ell)\}} |\theta_j|\} \max\{1, \sup_{x \in \mathbb{R}} |\mathbb{A}'(x)|\}]^L.
\end{aligned} \tag{22}$$

Proof of Corollary 2.4. Note that Lemma 2.3 proves (22). The proof of Corollary 2.4 is thus complete. \square

²Note that for all $d \in \mathbb{N}$, $v = (v_1, \dots, v_d) \in \mathbb{R}^d$ it holds that $\|v\| = [\sum_{i=1}^d |v_i|^2]^{1/2}$ (standard norm).

3. Error estimates for ANN approximations

In this section we establish in Corollary 3.7 below basic approximation error estimates for deep ANN approximations of twice continuously differentiable functions. In Section 4 we employ Corollary 3.7 together with Lemma 2.2 and Corollary 2.4 from Section 2 to establish in Proposition 4.5, Theorem 4.6, Corollary 4.7, and Corollary 4.8 error estimates for the deep Kolmogorov method and, thereby, prove Theorem 1.2 in the introduction. In our proof of Corollary 3.7 we also employ the well-known fact in Lemma 3.3 below that a constant function on a compact can be extended to a smooth function with the support in an open set containing the compact set. Only for completeness we include here a detailed proof for Lemma 3.3. Our proof of the ANN approximation result with convergence rates in Corollary 3.7 builds up on the findings in [79]. We also refer, for example, to [3, 54, 55, 77, 89, 103] and the references therein for further ANN approximation results.

3.1. Error estimates for shallow ANN approximations for C^2 -functions

Lemma 3.1. *Let $d \in \mathbb{N}$, $u \in C^2([-1, 1]^d, \mathbb{R})$. Then there exists $c \in (0, \infty)$ such that for all $\ell = (\ell_0, \ell_1, \ell_2) \in (\{d\} \times \mathbb{N} \times \{1\})$ there exists $\theta \in (-c, c)^{\mathfrak{d}(\ell)}$ such that*

$$\sup_{x \in (-1, 1)^d} |u(x) - \mathcal{N}_2^{\ell, \theta}(x)| \leq c(\ell_1)^{-\frac{1}{d+5}} \quad (23)$$

(cf. Definition 1.1).

Proof of Lemma 3.1. Throughout this proof for every $\ell = (\ell_0, \ell_1, \ell_2) \in \mathbb{N}^3$ let $N_\ell, \mathfrak{L}_\ell \in \mathbb{Z}$ satisfy

$$N_\ell = \lfloor (\ell_1)^{\frac{d+2}{d(d+4)}} \rfloor \quad \text{and} \quad \mathfrak{L}_\ell = \lceil \log_2(N_\ell) \rceil, \quad (24)$$

let $K_\ell: \mathbb{R} \rightarrow \mathbb{R}$ satisfy for all $t \in \mathbb{R}$ that

$$K_\ell(t) = \left[\int_{-\pi}^{\pi} \left(\frac{\sin(\frac{(N_\ell+1)s}{2})}{\sin(\frac{s}{2})} \right)^2 ds \right]^{-1} \left(\frac{\sin(\frac{(N_\ell+1)t}{2})}{\sin(\frac{t}{2})} \right)^2, \quad (25)$$

let $(\mathbf{a}_k^\ell)_{k \in \mathbb{Z}} \subseteq \mathbb{Z}$ satisfy for all $t \in \mathbb{R}$ that

$$K_\ell(t) = \sum_{k \in \mathbb{Z}} \mathbf{a}_k^\ell \exp(\mathbf{i}kt), \quad (26)$$

let $(a_k^\ell)_{k \in \mathbb{Z}^d} \subseteq \mathbb{Z}$ satisfy for all $k = (k_1, \dots, k_d) \in \mathbb{Z}^d$ that

$$a_k^\ell = \prod_{j=1}^d \mathbf{a}_{k_j}^\ell, \quad (27)$$

let $G_\ell: \mathbb{R}^d \rightarrow \mathbb{C}$ satisfy for all $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ that

$$G_\ell(x) = \sum_{k=(k_1, \dots, k_d) \in \mathbb{Z}^d} a_k^\ell \exp(\mathbf{i} \sum_{i=1}^d k_i x_i), \quad (28)$$

and let $J_\ell: C([- \pi, \pi]^d, \mathbb{R}) \times \mathbb{R}^d \rightarrow \mathbb{C}$ satisfy for all $g \in C([- \pi, \pi]^d, \mathbb{R})$, $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ that

$$J_\ell(g, x) = \int_{[- \pi, \pi]^d} G_\ell(x - y)g(y) dy. \quad (29)$$

Observe that [95, Chapter 6] demonstrates that there exist $F \in C^1([- \pi, \pi]^d, \mathbb{R})$, $\mathfrak{C} \in \mathbb{R}$ which satisfy that

- (i) it holds for all $x \in (-1, 1)^d$ that $F(x) = u(x)$,
- (ii) it holds for all $i \in \{1, 2, \dots, d\}$, $x = (x_1, \dots, x_d) \in [-\pi, \pi]^d$ with $x_i = -\pi$ that $F(x) = F(x_1, x_2, \dots, x_{i-1}, x_i + 2\pi, x_{i+1}, \dots, x_d)$, and
- (iii) it holds that $\sup_{x \in [-\pi, \pi]^d} |F(x)| \leq \mathfrak{C} \sup_{x \in (-1, 1)^d} |u(x)|$.

In the following for every $\ell \in \mathbb{N}^3$, $k = (k_1, \dots, k_d) \in \mathbb{Z}^d$ let $\mathfrak{J}_\ell(k) \in \mathbb{C}$ satisfy

$$\mathfrak{J}_\ell(k) = \frac{1}{(2\pi)^d} \int_{[-\pi, \pi]^d} J_\ell(F, y_1, y_2, \dots, y_d) \exp(-\mathbf{i} \sum_{i=1}^d k_i y_i) dy_1 dy_2 \dots dy_d \quad (30)$$

and for every $\ell \in \mathbb{N}^3$ let $v_\ell \in [0, \infty]$ satisfy

$$v_\ell = \sum_{k=(k_1, \dots, k_d) \in \mathbb{Z}^d} |\mathfrak{J}_\ell(k)| |k_1 + k_2 + \dots + k_d|^2. \quad (31)$$

Note that (3.8) in [79], (24), (25), and (29) establish that there exists $\rho \in \mathbb{R}$ which satisfies for all $\ell \in (\{d\} \times \mathbb{N} \times \{1\})$, $g \in C^1([-\pi, \pi]^d, \mathbb{R})$ that

$$\sup_{t \in [-\pi, \pi]^d} |J_\ell(g, t) - g(t)| \leq \rho(N_\ell)^{-1} \left[\sup_{x \in [-\pi, \pi]^d} \|(\nabla g)(x)\| \right]. \quad (32)$$

Observe that [79, Lemma 1] and (31) ensure that there exist $\theta_\ell = (\theta_\ell^1, \dots, \theta_\ell^{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $\ell \in (\{d\} \times \mathbb{N} \times \{1\})$, and $\varrho \in \mathbb{R}$ which satisfy for all $\ell = (\ell_0, \ell_1, \ell_2) \in (\{d\} \times \mathbb{N} \times \{1\})$ that

$$\sup_{i \in \{1, 2, \dots, \mathfrak{d}(\ell)\}} |\theta_\ell^i| \leq \max \left\{ 1, \frac{8\pi^2 v_\ell}{\ell_1} \right\} \quad (33)$$

$$\text{and} \quad \sup_{x \in (-1, 1)^d} |J_\ell(u, x) - \mathcal{N}_2^{\ell, \theta_\ell}(x)| \leq \varrho v_\ell \max\{\sqrt{\ln(\ell_1)}, \sqrt{d}\} (\ell_1)^{-\frac{1}{2} - \frac{1}{d}}. \quad (34)$$

(cf. Definition 1.1). Note that (24), (32), (34), and item (i) imply that for all $\ell = (\ell_0, \ell_1, \ell_2) \in (\{d\} \times \mathbb{N} \times \{1\})$ it holds that

$$\begin{aligned} & \sup_{x \in (-1, 1)^d} |u(x) - \mathcal{N}_2^{\ell, \theta_\ell}(x)| \\ & \leq \rho(N_\ell)^{-1} \sup_{x \in (-1, 1)^d} \|(\nabla u)(x)\| + \varrho v_\ell \max\{\sqrt{\ln(\ell_1)}, \sqrt{d}\} (\ell_1)^{-\frac{1}{2} - \frac{1}{d}} \\ & \leq \rho(\ell_1)^{-\frac{d+2}{d(d+4)}} \sup_{x \in (-1, 1)^d} \|(\nabla u)(x)\| + \varrho v_\ell \max\{\sqrt{\ln(\ell_1)}, \sqrt{d}\} (\ell_1)^{-\frac{1}{2} - \frac{1}{d}}. \end{aligned} \quad (35)$$

Moreover, observe that (4.1) in [79], (4.2) in [79], (4.6) in [79], and (24) show that there exist $c_1, c_2 \in \mathbb{R}$ such that for all $\ell \in (\{d\} \times \mathbb{N} \times \{1\})$ it holds that

$$v_\ell \leq 2d(4\pi c_1)^d c_2 \left[\sup_{x \in [-\pi, \pi]^d} \max\{|F(x)|, |(\nabla F)(x)|\} \right] 2^{-1} 3^{d/2} \sum_{i=0}^{\mathfrak{L}_\ell} (i+1)^d (2^i)^{1+\frac{d}{2}}. \quad (36)$$

The fact that $F \in C^1([-\pi, \pi]^d, \mathbb{R})$ hence proves that there exists $c \in \mathbb{R}$ such that for all $\ell \in (\{d\} \times \mathbb{N} \times \{1\})$ it holds that

$$v_\ell \leq c \sum_{i=0}^{\mathfrak{L}_\ell} (i+1)^d (2^i)^{1+\frac{d}{2}}. \quad (37)$$

Furthermore, note that (24) demonstrates that for all $\ell \in (\{d\} \times \mathbb{N} \times \{1\})$ it holds that

$$\begin{aligned} & \sum_{i=0}^{\mathfrak{L}_\ell} (i+1)^d (2^i)^{1+\frac{d}{2}} \leq (\mathfrak{L}_\ell + 1)^d \sum_{i=0}^{\mathfrak{L}_\ell} (2^i)^{1+\frac{d}{2}} = (\mathfrak{L}_\ell + 1)^d \frac{2^{1+\frac{d}{2}} (2^{1+\frac{d}{2}})^{\mathfrak{L}_\ell+1} - 1}{2^{1+\frac{d}{2}} - 1} \\ & \leq (\log_2(N_\ell) + 2)^d \frac{\sqrt{2}}{\sqrt{2}-1} (2^{1+\frac{d}{2}})^{L_\ell} \leq \frac{\sqrt{2}}{\sqrt{2}-1} (\log_2(N_\ell) + 2)^d (2N_\ell)^{1+\frac{d}{2}}. \end{aligned} \quad (38)$$

This, (24), and (37) establish that there exist $c_1, c_2 \in \mathbb{R}$ such that for all $\ell = (\ell_0, \ell_1, \ell_2) \in (\{d\} \times \mathbb{N} \times \{1\})$ it holds that

$$\begin{aligned} v_\ell &\leq c_1(\log_2(N_\ell) + 2)^d(2N_\ell)^{1+\frac{d}{2}} \\ &\leq c_1\left(\frac{d+2}{d(d+4)}\log_2(\ell_1) + 2\right)^d 2^{1+\frac{d}{2}}(\ell_1)^{\frac{(d+2)^2}{2d(d+4)}} \\ &\leq c_2(\ell_1)^{\frac{2}{d(d+4)}}(\ell_1)^{\frac{(d+2)^2}{2d(d+4)}} = c_2(\ell_1)^{\frac{1}{2}+\frac{4}{d(d+4)}}. \end{aligned} \quad (39)$$

Combining (33), (35), and the assumption that $u \in C^2([-1, 1]^d, \mathbb{R})$ therefore ensures that there exist $c_1, c_2, c_3 \in \mathbb{R}$ such that for all $\ell = (\ell_0, \ell_1, \ell_2) \in (\{d\} \times \mathbb{N} \times \{1\})$ it holds that

$$\begin{aligned} \sup_{x \in (-1, 1)^d} |u(x) - \mathcal{N}_2^{\ell, \theta_\ell}(x)| &\leq c_1(\ell_1)^{-\frac{d+2}{d(d+4)}} \sup_{x \in (-1, 1)^d} \|(\nabla u)(x)\| + c_1\sqrt{\ln(\ell_1)}(\ell_1)^{-\frac{1}{d+4}} \\ &\leq c_2(\ell_1)^{-\frac{d+2}{d(d+4)}} + c_2\sqrt{\ln(\ell_1)}(\ell_1)^{-\frac{1}{d+4}} \leq c_3(\ell_1)^{-\frac{1}{d+5}} \end{aligned} \quad (40)$$

$$\text{and} \quad \sup_{i \in \{1, 2, \dots, \mathfrak{d}(\ell)\}} |\theta_\ell^i| \leq \max\left\{1, c_3(\ell_1)^{-\frac{1}{2}+\frac{2}{d(d+4)}}\right\} \leq \max\{1, c_3\}. \quad (41)$$

Hence, we obtain that there exist $c_1, c_2 \in (0, \infty)$ such that for all $\ell = (\ell_0, \ell_1, \ell_2) \in (\{d\} \times \mathbb{N} \times \{1\})$ there exists $\theta \in (-c_2, c_2)^{\mathfrak{d}(\ell)}$ such that

$$\sup_{x \in (-1, 1)^d} |u(x) - \mathcal{N}_2^{\ell, \theta}(x)| \leq c_1(\ell_1)^{-\frac{1}{d+5}}. \quad (42)$$

This implies (23). The proof of Lemma 3.1 is thus complete. \square

Lemma 3.2. *Let $d \in \mathbb{N}$, $a \in \mathbb{R}$, $b \in (a, \infty)$, $u \in C^2([a, b]^d, \mathbb{R})$. Then there exists $c \in (0, \infty)$ such that for all $\ell = (\ell_0, \ell_1, \ell_2) \in (\{d\} \times \mathbb{N} \times \{1\})$ there exists $\theta \in (-c, c)^{\mathfrak{d}(\ell)}$ such that*

$$\sup_{x \in (a, b)^d} |u(x) - \mathcal{N}_2^{\ell, \theta}(x)| \leq c(\ell_1)^{-\frac{1}{d+5}} \quad (43)$$

(cf. Definition 1.1).

Proof of Lemma 3.2. Throughout this proof let $v: [-1, 1]^d \rightarrow \mathbb{R}$ satisfy for all $x \in [-1, 1]^d$ that

$$v(x) = u\left(\frac{b-a}{2}x + \left(\frac{a+b}{2}, \frac{a+b}{2}, \dots, \frac{a+b}{2}\right)\right). \quad (44)$$

Observe that Lemma 3.1 shows that there exist $c \in (0, \infty)$ and $\theta_\ell = (\theta_\ell^1, \dots, \theta_\ell^{\mathfrak{d}(\ell)}) \in (-c, c)^{\mathfrak{d}(\ell)}$, $\ell \in \mathbb{N}^3$, which satisfy for all $\ell = (\ell_0, \ell_1, \ell_2) \in (\{d\} \times \mathbb{N} \times \{1\})$ that

$$\sup_{x \in (-1, 1)^d} |v(x) - \mathcal{N}_2^{\ell, \theta_\ell}(x)| \leq c(\ell_1)^{-\frac{1}{d+5}} \quad (45)$$

(cf. Definition 1.1). In the following for every $\ell = (\ell_0, \ell_1, \ell_2) \in \mathbb{N}^3$ let $\vartheta_\ell = (\vartheta_\ell^1, \dots, \vartheta_\ell^{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$ satisfy that

- (i) it holds for all $i \in \{1, 2, \dots, \ell_1 \ell_0\}$ that $\vartheta_\ell^i = \frac{2}{b-a}\theta_\ell^i$,
- (ii) it holds for all $i \in \{\ell_1 \ell_0 + 1, \ell_1 \ell_0 + 2, \dots, \ell_1 \ell_0 + \ell_1\}$ that $\vartheta_\ell^i = \theta_\ell^i - \frac{a+b}{2}$, and
- (iii) it holds for all $i \in \{\ell_1 \ell_0 + \ell_1 + 1, \ell_1 \ell_0 + \ell_1 + 2, \dots, \mathfrak{d}(\ell)\}$ that $\vartheta_\ell^i = \theta_\ell^i$.

Note that item (i), item (ii), and item (iii) prove that for all $\ell \in (\{d\} \times \mathbb{N} \times \{1\})$ it holds that

$$\mathcal{N}_2^{\ell, \theta_\ell}(x) = \mathcal{N}_2^{\ell, \vartheta_\ell}\left(\frac{b-a}{2}x + \left(\frac{a+b}{2}, \frac{a+b}{2}, \dots, \frac{a+b}{2}\right)\right). \quad (46)$$

Combining this, (44), and (45) demonstrates for all $\ell \in (\{d\} \times \mathbb{N} \times \{1\})$ that

$$\sup_{x \in (-1,1)^d} \left| u\left(\frac{b-a}{2}x + \left(\frac{a+b}{2}, \frac{a+b}{2}, \dots, \frac{a+b}{2}\right)\right) - \mathcal{N}_2^{\ell, \vartheta_\ell}\left(\frac{b-a}{2}x + \left(\frac{a+b}{2}, \frac{a+b}{2}, \dots, \frac{a+b}{2}\right)\right) \right| \leq (\ell_1)^{\frac{-1}{d+5}}. \quad (47)$$

Hence, we obtain for all $\ell \in (\{d\} \times \mathbb{N} \times \{1\})$ that

$$\sup_{x \in (a,b)^d} |u(x) - \mathcal{N}_2^{\ell, \vartheta_\ell}(x)| \leq c(\ell_1)^{\frac{-1}{d+5}}. \quad (48)$$

In the next step we combine (45), item (i), item (ii), and item (iii) to obtain for all $\ell \in (\{d\} \times \mathbb{N} \times \{1\})$, $i \in \{1, 2, \dots, \mathfrak{d}(\ell)\}$ that

$$|\vartheta_\ell^i| \leq c(1 + |\frac{b-a}{2}|) + |\frac{a+b}{2}|. \quad (49)$$

This and (48) establish (43). The proof of Lemma 3.2 is thus complete. \square

3.2. Error estimates for deep ANN approximations for C^2 -functions

Lemma 3.3. *Let $d \in \mathbb{N}$, let $O \subseteq \mathbb{R}^d$ be open, and let $K \subseteq O$ be compact. Then there exists $\phi \in C^\infty(\mathbb{R}^d, \mathbb{R})$ such that*

(i) *it holds for all $x \in K$ that $\phi(x) = 1$ and*

(ii) *it holds for all $x \in \mathbb{R}^d \setminus O$ that $\phi(x) = 0$.*

Proof of Lemma 3.3. Throughout this proof assume without loss of generality that $K \neq \emptyset$ (otherwise we can choose $\phi = (\mathbb{R}^d \ni x \mapsto 0 \in \mathbb{R})$), for every $r \in \mathbb{R}$, $x \in \mathbb{R}^d$ let $B_r(x) \subseteq \mathbb{R}^d$ satisfy $B_r(x) = \{y \in \mathbb{R}^d : \|x - y\| < r\}$, let $R \in (0, \infty)$ satisfy $K \subseteq B_R(0)$, let $\mathbb{O} \subseteq \mathbb{R}^d$ satisfy $\mathbb{O} = O \cap B_R(0)$, let $r \in \mathbb{R}$ satisfy

$$r = \frac{1}{3} \left[\inf_{x \in K} \inf_{y \in (\mathbb{R}^d \setminus \mathbb{O})} \|x - y\| \right], \quad (50)$$

for every $x \in \mathbb{R}^d$ let $f_x : \mathbb{R}^d \rightarrow \mathbb{R}$ satisfy for all $y \in \mathbb{R}^d$ that

$$f_x(y) = \begin{cases} \exp(\left[\|x - y\|^2 - r^2 \right]^{-1}) & : \|x - y\| < r, \\ 0 & : \|x - y\| \geq r, \end{cases} \quad (51)$$

let $N \in \mathbb{N} \cap (r^{-1}2d^{1/2}, \infty)$, let $I \subseteq \mathbb{R}^d$ satisfy

$$I = \{v \in \mathbb{R}^d : Nv \in \mathbb{Z}^d\}, \quad (52)$$

and let $\phi : \mathbb{R}^d \rightarrow \mathbb{R}$ satisfy for all $x \in \mathbb{R}^d$ that

$$\phi(x) = \left(\sum_{v \in I, \text{supp}(f_v) \cap K \neq \emptyset} f_v(x) \right) \left(\sum_{v \in I} f_v(x) \right)^{-1} \quad (53)$$

Observe that (51), the fact that $([0, \infty) \ni x \mapsto \exp(\frac{1}{x-r^2+\mathbb{1}_{[0,r^2]}(x)}) \mathbb{1}_{[0,r^2]}(x) \in \mathbb{R}) \in C^\infty([0, \infty), \mathbb{R})$, and the fact that for all $v \in \mathbb{R}^d$ it holds that $(\mathbb{R}^d \ni w \mapsto \|v - w\|^2 \in \mathbb{R}) \in C^\infty(\mathbb{R}^d, \mathbb{R})$ ensure that for all $v \in \mathbb{R}^d$ it holds that

$$f_v \in C^\infty(\mathbb{R}^d, \mathbb{R}). \quad (54)$$

Next, note that (51) and (52) imply for all $s \in (0, \infty)$, $x \in B_s(0)$ that

$$\sum_{v \in I} f_v(x) = \sum_{v \in I \cap B_{r+s}(0)} f_v(x) \quad (55)$$

$$\text{and} \quad \sum_{v \in I, \text{supp}(f_v) \cap K \neq \emptyset} f_v(x) = \sum_{v \in I \cap B_{r+s}(0), \text{supp}(f_v) \cap K \neq \emptyset} f_v(x). \quad (56)$$

In addition, observe that (52) shows for all $s \in (0, \infty)$ that

$$\#\{I \cap B_s(0)\} < \infty. \quad (57)$$

Combining this and (55) proves for all $s \in (0, \infty)$ that

$$(B_s(0) \ni x \mapsto \sum_{v \in I} f_v(x) \in \mathbb{R}) \in C^\infty(B_s(0), \mathbb{R}) \quad (58)$$

$$\text{and} \quad (B_s(0) \ni x \mapsto \sum_{v \in I, \text{supp}(f_v) \cap K \neq \emptyset} f_v(x) \in \mathbb{R}) \in C^\infty(B_s(0), \mathbb{R}). \quad (59)$$

Hence, we obtain that

$$(\mathbb{R}^d \ni x \mapsto \sum_{v \in I} f_v(x) \in \mathbb{R}) \in C^\infty(\mathbb{R}^d, \mathbb{R}) \quad (60)$$

$$\text{and} \quad (\mathbb{R}^d \ni x \mapsto \sum_{v \in I, \text{supp}(f_v) \cap K \neq \emptyset} f_v(x) \in \mathbb{R}) \in C^\infty(\mathbb{R}^d, \mathbb{R}). \quad (61)$$

Moreover, note that the fact that $N \geq \frac{2\sqrt{d}}{r}$ demonstrates that for all $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ there exists $y = (y_1, y_2, \dots, y_d) \in \mathbb{R}^d$ such that for all $i \in \{1, 2, \dots, d\}$ it holds that

$$|x_i - y_i| < \frac{r}{\sqrt{d}} \quad \text{and} \quad Ny_i \in \mathbb{Z}. \quad (62)$$

This and (52) establish for all $x \in \mathbb{R}^d$ that there exists $y \in I$ such that $\|x - y\| < r$. Combining this and (51) ensures for all $x \in \mathbb{R}^d$ it holds that

$$\sum_{v \in I} f_v(x) > 0. \quad (63)$$

This, (53), (60), and (61) imply that

$$\phi \in C^\infty(\mathbb{R}^d, \mathbb{R}). \quad (64)$$

Furthermore, observe that (51) shows that for all $v \in \mathbb{R}^d$ it holds that

$$\text{supp}(f_v) = \overline{B_r(v)}. \quad (65)$$

Hence, we obtain for all $x \in K$, $v \in I \cap \overline{B_r(x)}$ that

$$\text{supp}(f_v) \cap K \neq \emptyset. \quad (66)$$

Combining this, (53), and the fact that for all $x \in \mathbb{R}^d$ it holds that $\sum_{v \in I} f_v(x) = \sum_{v \in I \cap \overline{B_r(x)}} f_v(x)$ proves that for all $x \in K$ it holds that

$$\phi(x) = \left(\sum_{v \in I \cap \overline{B_r(x)}} f_v(x) \right) \left(\sum_{v \in I \cap \overline{B_r(x)}} f_v(x) \right)^{-1} = 1. \quad (67)$$

Next we combine (50) and (65) to obtain for all $x \in \mathbb{R}^d \setminus \mathbb{O}$, $v \in \overline{B_r(x)}$ it holds that

$$\text{supp}(f_v) \cap K = \overline{B_r(v)} \cap K \subseteq \overline{B_{2r}(x)} \cap K = \emptyset. \quad (68)$$

The fact that for all $x \in \mathbb{R}^d$ it holds that $\sum_{v \in I, \text{supp}(f_v) \cap K \neq \emptyset} f_v(x) = \sum_{v \in I \cap \overline{B_r(x)}, \text{supp}(f_v) \cap K \neq \emptyset} f_v(x)$ hence demonstrates for all $x \in \mathbb{R}^d \setminus \mathbb{O}$ that

$$\phi(x) = \left(\sum_{v \in I \cap \overline{B_r(x)}, \text{supp}(f_v) \cap K \neq \emptyset} f_v(x) \right) \left(\sum_{v \in I} f_v(x) \right)^{-1} = 0. \quad (69)$$

This, (64), and (67) establish items (i) and item (ii). The proof of Lemma 3.3 is thus complete. \square

Lemma 3.4. *Let $d \in \mathbb{N}$, let $D \subseteq \mathbb{R}^d$ be bounded, and let $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d\} \times \mathbb{N}^{L-1} \times \{1\})$, $\mathbf{1} \in (\{d\} \times \mathbb{N} \times \{1\})$, $\vartheta \in \mathbb{R}^{\mathfrak{d}(\mathbf{1})}$. Then there exists $\theta \in [-\|\vartheta\| - 1, \|\vartheta\| + 1]^{\mathfrak{d}(\ell)}$ which satisfies for all $x \in D$ that*

$$\mathcal{N}_L^{\ell, \theta}(x) = \mathcal{N}_2^{\mathbf{1}, \vartheta}(x). \quad (70)$$

Proof of Lemma 3.4. Throughout this proof let $\xi = (\xi_1, \dots, \xi_{\mathfrak{d}(\mathbf{1})}) \in \mathbb{R}^{\mathfrak{d}(\mathbf{1})}$ satisfy $\xi = \vartheta$. Note that, for instance, [67, Lemma 4.4.7] (applied with $2 \curvearrowright L$, $\mathfrak{L} \curvearrowright L$, $l \curvearrowright \mathbf{1}$, $\mathfrak{l} \curvearrowright \ell$ in the notation of [67, Lemma 4.4.7]) ensures that there exists $\theta = (\theta_1, \dots, \theta_{\mathfrak{d}(\ell)}) \in \mathbb{R}^{\mathfrak{d}(\ell)}$ which satisfies that

(i) it holds for all $x \in \mathbb{R}^{\mathfrak{d}}$ that $\mathcal{N}_L^{\ell, \theta}(x) = \mathcal{N}_2^{\mathbf{1}, \xi}(x)$ and

(ii) it holds that $\max\{|\theta_1|, |\theta_2|, \dots, |\theta_{\mathfrak{d}(\ell)}|\} \leq \max\{1, \max\{|\xi_1|, |\xi_2|, \dots, |\xi_{\mathfrak{d}(\mathbf{1})}|\}\}$.

(cf. Definition 1.1). Observe that item (i), item (ii), and the fact that $\max\{1, \max\{|\xi_1|, |\xi_2|, \dots, |\xi_{\mathfrak{d}(\mathbf{1})}|\}\} \leq \|\xi\| + 1$ imply (70). The proof of Lemma 3.4 is thus complete. \square

Lemma 3.5. *Let $d \in \mathbb{N}$, $a \in \mathbb{R}$, $b \in (a, \infty)$, $u \in C^2([a, b]^d, \mathbb{R})$. Then there exists $c \in (0, \infty)$ such that for all $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d\} \times \mathbb{N}^{L-1} \times \{1\})$ there exists $\theta \in (-c, c)^{\mathfrak{d}(\ell)}$ such that*

$$\int_{(a,b)^d} |u(x) - \mathcal{N}_L^{\ell, \theta}(x)|^2 dx \leq c[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{\frac{-2}{d+5}} \quad (71)$$

(cf. Definition 1.1).

Proof of Lemma 3.5. Throughout this proof let $\mathbf{1} \in \mathbb{N}^3$, $\vartheta \in \mathbb{R}^{\mathfrak{d}(\mathbf{1})}$ satisfy $\mathbf{1} = (1, 2, 1)$ and $\vartheta = (1, -1, 0, 0, 1, -1, 0)$. Note that, for example, [67, Lemma 2.2.7] shows that for all $x \in \mathbb{R}$ it holds that

$$\mathcal{N}_2^{\mathbf{1}, \vartheta}(x) = x. \quad (72)$$

(cf. Definition 1.1). Observe that Lemma 3.1 proves that there exist $c \in (0, \infty)$ and $\eta: (\{d\} \times (\cup_{L=0}^{\infty} \mathbb{N}^L) \times \{1\}) \rightarrow (-c, c)^{\mathfrak{d}(\ell)}$ which satisfy for all $\ell = (\ell_0, \ell_1, \ell_2) \in (\{d\} \times \mathbb{N} \times \{1\})$ that

$$\int_{(a,b)^d} |u(x) - \mathcal{N}_2^{\ell, \eta(\ell)}(x)|^2 dx \leq c(\ell_1)^{\frac{-2}{d+5}}. \quad (73)$$

Note that (1) and the fact that u is bounded demonstrate that there exists $C \in \mathbb{R}$ such that for all $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d\} \times (\mathbb{N}^{L-1} \times \{1\}))$ with $\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\} = 1$ it holds that

$$\int_{(a,b)^d} |u(x) - \mathcal{N}_L^{\ell, 0}(x)|^2 dx = \int_{(a,b)^d} |u(x)|^2 dx \leq C = C[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{\frac{-2}{d+5}}. \quad (74)$$

In the next step we observe that Lemma 3.4 and (72) establish that for all $\ell = (\ell_0, \ell_1, \ell_2) \in (\{d\} \times \mathbb{N} \times \{1\})$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d\} \times (\mathbb{N} \cap [\max\{2, \ell_1\}, \infty))^{L-1} \times \{1\})$ there exists $\theta \in [-\|\vartheta\| - 1, \|\vartheta\| + 1]^{\mathfrak{d}(\ell)}$ such that for all $x \in \mathbb{R}^d$ it holds that

$$\mathcal{N}_L^{\ell, \theta}(x) = \mathcal{N}_2^{\ell, \eta(\ell)}(x). \quad (75)$$

Hence, we obtain that for all $\ell = (\ell_0, \ell_1, \ell_2) \in (\{d\} \times (\mathbb{N} \cap [2, \infty)) \times \{1\})$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d\} \times (\mathbb{N} \cap [\ell_1, \infty))^{L-1} \times \{1\})$ there exists $\theta \in (-\max\{\|\vartheta\| + 1, c\}, \max\{\|\vartheta\| + 1, c\})^{\mathfrak{d}(\ell)}$ such that for all $x \in \mathbb{R}^d$ it holds that

$$\mathcal{N}_L^{\ell, \theta}(x) = \mathcal{N}_2^{\ell, \eta(\ell)}(x). \quad (76)$$

Combining this and (73) ensures that for all $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d\} \times \mathbb{N}^{L-1} \times \{1\})$ with $\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\} \geq 2$ there exists $\theta \in (-\max\{\|\vartheta\| + 1, c\}, \max\{\|\vartheta\| + 1, c\})^{\mathfrak{d}(\ell)}$ such that

$$\int_{(a,b)^d} |u(x) - \mathcal{N}_L^{\ell, \theta}(x)|^2 dx \leq c[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{\frac{-2}{d+5}}. \quad (77)$$

This and (74) imply (71). The proof of Lemma 3.5 is thus complete. \square

Corollary 3.6. *Let $d \in \mathbb{N}$, let $D \subseteq \mathbb{R}^d$ be measurable and bounded, and³ let $u \in C^2(\bar{D}, \mathbb{R})$. Then there exists $c \in (0, \infty)$ such that for all $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d\} \times \mathbb{N}^{L-1} \times \{1\})$ there exists $\theta \in (-c, c)^{\mathfrak{d}(\ell)}$ such that*

$$\int_D |u(x) - \mathcal{N}_L^{\ell, \theta}(x)|^2 dx \leq c[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{\frac{-2}{\bar{d}+5}} \quad (78)$$

(cf. Definition 1.1).

Proof of Corollary 3.6. Note that the assumption that D is bounded shows that there exist $a \in \mathbb{R}$, $b \in (a, \infty)$ which satisfy

$$\bar{D} \subseteq (a, b)^d. \quad (79)$$

Observe that the assumption that $u \in C^2(\bar{D}, \mathbb{R})$ proves that there exists an open set $V \subseteq \mathbb{R}^d$ and $\bar{u} \in C^2(V, \mathbb{R})$ which satisfy for all $x \in \bar{D}$ that

$$\bar{D} \subseteq V \quad \text{and} \quad u(x) = \bar{u}(x). \quad (80)$$

Note that (80), Lemma 3.3, and the assumption that D is bounded demonstrate that there exists $\phi \in C^\infty(\mathbb{R}^d, \mathbb{R})$ which satisfies for all $x \in \bar{D}$, $y \in \mathbb{R}^d \setminus V$ that

$$\phi(x) = 1 \quad \text{and} \quad \phi(y) = 0. \quad (81)$$

In the following let $v: \mathbb{R}^d \rightarrow \mathbb{R}$ satisfy for all $x \in \mathbb{R}^d$ that

$$v(x) = \begin{cases} \phi(x)\bar{u}(x) & : x \in V \\ 0 & : x \in \mathbb{R}^d \setminus V. \end{cases} \quad (82)$$

Observe that (80), (81), and (82) establish that for all $x \in \bar{D}$ it holds that

$$v(x) = \phi(x)\bar{u}(x) = \phi(x)u(x) = u(x). \quad (83)$$

Next, note that (81), (82), and the fact that $\bar{u} \in C^2(V, \mathbb{R})$ ensure that $v \in C^1(\mathbb{R}^d, \mathbb{R})$. Combining this, (79), (83), and Lemma 3.5 implies that there exists $c \in (0, \infty)$ such that for all $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d\} \times \mathbb{N}^{L-1} \times \{1\})$ there exists $\theta \in (-c, c)^{\mathfrak{d}(\ell)}$ such that

$$\begin{aligned} \int_D |u(x) - \mathcal{N}_L^{\ell, \theta}(x)|^2 dx &\leq \int_{(a,b)^d} |v(x) - \mathcal{N}_L^{\ell, \theta}(x)|^2 dx \\ &\leq c[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{\frac{-2}{\bar{d}+5}} \end{aligned} \quad (84)$$

(cf. Definition 1.1). This shows (78). The proof of Corollary 3.6 is thus complete. \square

Corollary 3.7. *Let $d \in \mathbb{N}$, let $D \subseteq \mathbb{R}^d$ be measurable and bounded, and let $u \in C^2(\bar{D}, \mathbb{R})$. Then there exists $c \in (0, \infty)$ such that for all $R \in (0, \infty]$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d\} \times \mathbb{N}^{L-1} \times \{1\})$ there exists $\theta \in (-\min\{c, R\}, \min\{c, R\})^{\mathfrak{d}(\ell)}$ such that*

$$\int_D |u(x) - \mathcal{N}_L^{\ell, \theta}(x)|^2 dx \leq c\mathbb{1}_{(-\infty, c)}(R) + c[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{\frac{-2}{\bar{d}+5}} \quad (85)$$

(cf. Definition 1.1).

³Note that for every $d, n \in \mathbb{N}$ and every closed $D \subseteq \mathbb{R}^d$ it holds that $C^n(D, \mathbb{R}) = \{f \in C(D, \mathbb{R}) : (\exists g \in C^n(\mathbb{R}^d, \mathbb{R}) : g|_D = f)\}$.

Proof of Corollary 3.7. Observe that Corollary 3.6 proves that there exists $c \in (0, \infty)$ such that for all $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d\} \times \mathbb{N}^{L-1} \times \{1\})$ there exists $\theta \in (-c, c)^{\mathfrak{d}(\ell)}$ such that

$$\int_D |u(x) - \mathcal{N}_L^{\ell, \theta}(x)|^2 dx \leq c[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{\frac{-2}{d+5}} \quad (86)$$

(cf. Definition 1.1). Note that (86) demonstrates that for all $R \in [c, \infty]$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d\} \times \mathbb{N}^{L-1} \times \{1\})$ there exists $\theta \in (-\min\{c, R\}, \min\{c, R\})^{\mathfrak{d}(\ell)}$ such that

$$\int_D |u(x) - \mathcal{N}_L^{\ell, \theta}(x)|^2 dx \leq c[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{\frac{-2}{d+5}}. \quad (87)$$

In addition, observe that (1), the fact that u is bounded, and the assumption that f is non-decreasing establish that there exists $\mathfrak{C} \in (0, \infty)$ such that for all $R \in (0, c)$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d\} \times \mathbb{N}^{L-1} \times \{1\})$ there exists $\theta \in (-\min\{c, R\}, \min\{c, R\})^{\mathfrak{d}(\ell)}$ such that

$$\int_D |u(x) - \mathcal{N}_L^{\ell, 0}(x)|^2 dx = \int_D |u(x)|^2 dx \leq \mathfrak{C} = \mathfrak{C} \mathbb{1}_{(-\infty, c)}(R). \quad (88)$$

Combining this and (87) ensures (85). The proof of Corollary 3.7 is thus complete. \square

4. Error estimates for the deep Kolmogorov method

In this section we employ Lemma 2.2 and Corollary 2.4 from Section 2 and Corollary 3.7 from Section 3 to establish in Proposition 4.5, Theorem 4.6, Corollary 4.7, and Corollary 4.8 below error estimates for the deep Kolmogorov method. Theorem 1.2 in the introduction is an immediate consequence of Corollary 4.8. Our proof of Proposition 4.5 also employs the well-known Sobolev estimates with explicit embedding constants in Lemma 4.1 and Corollary 4.2 below, the Feynman-Kac representation for solutions of heat PDEs that we recall in Lemma 4.3 below, and the well-known error estimates for the Monte Carlo method in Lemma 4.4 below. We also refer, for instance, to [92, Lemma 4] and [93, Appendix C] for abstract general error estimates for ANN approximations in terms of the optimization error.

4.1. Sobolev embedding estimates with explicit constants

Lemma 4.1. *Let $d \in \mathbb{N}$, $\Phi \in C^1((0, 1)^d, \mathbb{R})$. Then*

$$\sup_{x \in (0, 1)^d} |\Phi(x)| \leq 8\sqrt{e} \left[\int_{(0, 1)^d} (|\Phi(x)|^{\max\{2, d^2\}} + \|(\nabla \Phi)(x)\|^{\max\{2, d^2\}}) dx \right]^{1/\max\{2, d^2\}}. \quad (89)$$

Proof of Lemma 4.1. Note that, for example, [47, Corollary 2.15] implies (89). The proof of Lemma 4.1 is thus complete. \square

Corollary 4.2. *Let $d \in \mathbb{N}$, $a \in \mathbb{R}$, $b \in (a, \infty)$, $p \in [\max\{2, d^2\}, \infty)$, $\Phi \in C^1((a, b)^d, \mathbb{R})$. Then*

$$\sup_{x \in (a, b)^d} |\Phi(x)| \leq 16\sqrt{e} \max\{(b-a)^{-d/p}, b-a\} \left[\int_{(a, b)^d} (|\Phi(x)|^p + \|(\nabla \Phi)(x)\|^p) dx \right]^{1/p}. \quad (90)$$

Proof of Corollary 4.2. Throughout this proof let $\mathbf{e} \in \mathbb{R}^d$ satisfy $\mathbf{e} = (1, 1, \dots, 1)$ and let $\Psi: (0, 1)^d \rightarrow \mathbb{R}$ satisfy for all $x = (x_1, \dots, x_d) \in (0, 1)^d$ that

$$\Psi(x) = \Phi(a\mathbf{e} + (b-a)x). \quad (91)$$

Observe that (91) and the assumption that $\Phi \in C^1((a, b)^d, \mathbb{R})$ show that $\Psi \in C^1((0, 1)^d, \mathbb{R})$. Lemma 4.1, the assumption that $p \geq \max\{2, d^2\}$, and Holder inequality therefore prove that

$$\begin{aligned} \sup_{x \in (0, 1)^d} |\Psi(x)| &\leq 8\sqrt{e} \left[\int_{(0, 1)^d} (|\Psi(x)|^{\max\{2, d^2\}} + \|(\nabla \Psi)(x)\|^{\max\{2, d^2\}}) dx \right]^{1/\max\{2, d^2\}} \\ &\leq 8\sqrt{e} \left[\int_{(0, 1)^d} (|\Psi(x)|^{\max\{2, d^2\}} + \|(\nabla \Psi)(x)\|^{\max\{2, d^2\}})^{p/\max\{2, d^2\}} dx \right]^{1/p}. \end{aligned} \quad (92)$$

This, the fact that for all $x, y \in \mathbb{R}^d$, $p \in [1, \infty)$ it holds that $\|x + y\|^p \leq 2^p \|x\|^p + 2^p \|y\|^p$, and the fact that $\max\{2, d^2\} \geq 1$ demonstrate that

$$\sup_{x \in (0, 1)^d} |\Psi(x)| \leq 16\sqrt{e} \left[\int_{(0, 1)^d} (|\Psi(x)|^p + \|(\nabla \Psi)(x)\|^p) dx \right]^{1/p}. \quad (93)$$

Moreover, note that (91) establishes that for all $x \in (0, 1)^d$ it holds that

$$(\nabla \Psi)(x) = (b - a)(\nabla \Phi)(a\mathbf{e} + (b - a)x) \quad (94)$$

Combining this and (91) ensures that

$$\begin{aligned} &\int_{(0, 1)^d} (|\Psi(x)|^p + \|(\nabla \Psi)(x)\|^p) dx \\ &\leq \max\{1, (b - a)^p\} \left[\int_{(0, 1)^d} (|\Psi(a\mathbf{e} + (b - a)x)|^p + \|(\nabla \Psi)(a\mathbf{e} + (b - a)x)\|^p) dx \right] \\ &= \frac{\max\{1, (b - a)^p\}}{(b - a)^d} \left[\int_{(0, 1)^d} (|\Psi(a\mathbf{e} + (b - a)x)|^p \right. \\ &\quad \left. + \|(\nabla \Psi)(a\mathbf{e} + (b - a)x)\|^p) d(a\mathbf{e} + (b - a)x) \right] \\ &= \frac{\max\{1, (b - a)^p\}}{(b - a)^d} \left[\int_{(a, b)^d} (|\Phi(x)|^p + \|(\nabla \Phi)(x)\|^p) dx \right] \\ &= \max\{(b - a)^{-d}, (b - a)^p\} \left[\int_{(a, b)^d} (|\Phi(x)|^p + \|(\nabla \Phi)(x)\|^p) dx \right]. \end{aligned} \quad (95)$$

This, (91), and (93) imply (90). The proof of Corollary 4.2 is thus complete. \square

4.2. Feynman-Kac representations for solutions of heat PDEs

Lemma 4.3. *Let $T \in (0, \infty)$, $d \in \mathbb{N}$, let $u \in C([0, T] \times \mathbb{R}^d, \mathbb{R})$ be at most polynomially growing, assume $u|_{(0, T) \times \mathbb{R}^d} \in C^{1,2}((0, T) \times \mathbb{R}^d, \mathbb{R})$, assume for all $t \in (0, T)$, $x \in \mathbb{R}^d$ that*

$$\frac{\partial}{\partial t} u(t, x) + \frac{1}{2} \Delta_x u(t, x) = 0, \quad (96)$$

let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $W : [0, T] \times \Omega \rightarrow \mathbb{R}^d$ be a standard Brownian motion, let $\mathbb{T} : \Omega \rightarrow [0, T]$ and $\mathbb{X} : \Omega \rightarrow \mathbb{R}^d$ be random variables, let $q \in [0, \infty)$ satisfy $\mathbb{E}[\|\mathbb{X}\|^{2q}] + \sup_{x \in \mathbb{R}^d} ((1 + |x|)^{-q} |u(T, x)|) < \infty$, and assume that (\mathbb{T}, \mathbb{X}) and W are independent. Then it holds \mathbb{P} -a.s. that

$$\mathbb{E}[u(T, \mathbb{X} + W_{T-\mathbb{T}}) | (\mathbb{T}, \mathbb{X})] = u(\mathbb{T}, \mathbb{X}). \quad (97)$$

Proof of Lemma 4.3. Throughout this proof let $g : \mathbb{R}^d \rightarrow \mathbb{R}$ satisfy for all $x \in \mathbb{R}^d$ that $g(x) = u(T, x)$, let $v : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ satisfy for all $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$v(t, x) = u(T - t, x). \quad (98)$$

Observe that (96) and (98) show that for all $t \in (0, T)$, $x \in \mathbb{R}^d$ it holds that

$$\frac{\partial}{\partial t} v(t, x) - \frac{1}{2} \Delta_x v(t, x) = 0 \quad \text{and} \quad v(0, x) = g(x), \quad (99)$$

Combining this and [57, Corollary 4.17] proves for all $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$v(t, x) = \mathbb{E}[g(x + W_t)]. \quad (100)$$

This and (98) demonstrate for all $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$u(t, x) = \mathbb{E}[g(x + W_{T-t})]. \quad (101)$$

In the following let $D = [0, T] \times \mathbb{R}^d$, $E = C([0, T], \mathbb{R}^d)$, let $\Phi: D \times E \rightarrow [-\infty, \infty]$ satisfy for all $(t, x) \in [0, T] \times \mathbb{R}^d$, $\mathbf{W} \in C([0, T] \times \mathbb{R}^d)$ that

$$\Phi(t, x, \mathbf{W}) = g(x, \mathbf{W}_{T-t}), \quad (102)$$

and let $\phi: D \rightarrow [-\infty, \infty]$ satisfy for all $(t, x) \in D$ that

$$\phi(t, x) = \mathbb{E}[\Phi(t, x, W)]. \quad (103)$$

Furthermore, note that (102), the assumption that $\mathbb{E}[\|\mathbb{X}\|^q] < \infty$, the assumption that $\sup_{x \in \mathbb{R}^d} ((1 + \|x\|)^{-q} |g(x)|) < \infty$, and the fact that $\mathbb{P}(|\mathbb{T}| \leq T) = 1$ the assumption that W is a standard Brownian motion establish that there exists $c \in \mathbb{R}$ such that

$$\mathbb{E}[|\Phi(\mathbb{T}, \mathbb{X}, W)|] = \mathbb{E}[|g(\mathbb{X}, W_{T-\mathbb{T}})|] \leq c(1 + \mathbb{E}[\|\mathbb{X}\|^q] + \mathbb{E}[\sup_{t \in [0, T]} \|W_t\|^q]) < \infty. \quad (104)$$

Next we combine (102), the assumption that $\sup_{x \in \mathbb{R}^d} ((1 + \|x\|)^{-q} |g(x)|) < \infty$, and the assumption that W is a standard Brownian motion to obtain that for all $(t, x) \in D$ there exists $c \in \mathbb{R}$ such that

$$\mathbb{E}[|\Phi(t, x, W)|] = \mathbb{E}[|g(x, W_{T-t})|] \leq c(1 + \mathbb{E}[\sup_{t \in [0, T]} \|W_t\|^q]) < \infty. \quad (105)$$

Combining this, (102), (103), (104), the assumption that (\mathbb{T}, \mathbb{X}) and W are independent, and [29, Proposition 3.15] (applied⁴ with $\Omega \curvearrowright \Omega$, $\mathcal{F} \curvearrowright \mathcal{F}$, $\mathbb{P} \curvearrowright \mathbb{P}$, $D \curvearrowright D$, $E \curvearrowright E$, $\mathcal{G} \curvearrowright \sigma((\mathbb{T}, \mathbb{X}))$, $X \curvearrowright (\mathbb{T}, \mathbb{X})$, $Y \curvearrowright W$, $\Phi \curvearrowright \Phi$, $\phi \curvearrowright \phi$ in the notation of [29, Proposition 3.15]) ensures that it holds \mathbb{P} -a.s. that

$$\mathbb{E}[g(\mathbb{X} + W_{T-\mathbb{T}}) | (\mathbb{T}, \mathbb{X})] = \mathbb{E}[\Phi(\mathbb{X}, \mathbb{T}, W) | (\mathbb{T}, \mathbb{X})] = \phi(\mathbb{T}, \mathbb{X}). \quad (106)$$

This and (101) imply that it holds \mathbb{P} -a.s. that

$$\mathbb{E}[g(\mathbb{X} + W_{T-\mathbb{T}}) | (\mathbb{T}, \mathbb{X})] = u(\mathbb{T}, \mathbb{X}). \quad (107)$$

This shows (97). The proof of Lemma 4.3 is thus complete. \square

4.3. Error estimates for the Monte Carlo method

Lemma 4.4. *Let $p \in [2, \infty)$, $d, n \in \mathbb{N}$, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, and let $X_i: \Omega \rightarrow \mathbb{R}^d$, $i \in \{1, 2, \dots, n\}$, be i.i.d. random variables with $\mathbb{E}[\|X_1\|] < \infty$. Then*

$$\left(\mathbb{E}[\|\mathbb{E}[X_1] - \frac{1}{n}(\sum_{i=1}^n X_i)\|^p] \right)^{1/p} \leq 2 \left[\frac{p-1}{n} \right]^{1/2} (\mathbb{E}[\|X_1 - \mathbb{E}[X_1]\|^p])^{1/p}. \quad (108)$$

Proof of Lemma 4.4. Observe that, for instance, [52, Corollary 2.6] proves (108). The proof of Lemma 4.4 is thus complete. \square

⁴Note that for every set Ω , every measurable space (S, \mathcal{S}) , and every function $X: \Omega \rightarrow S$ it holds that $\sigma(X) = \{A \subseteq \Omega: (\exists B \in \mathcal{S}: A = X^{-1}(B))\}$ (the smallest sigma-algebra on Ω with respect to which X is measurable).

4.4. Error estimates for the deep Kolmogorov method

Proposition 4.5. *Let $T, \kappa \in (0, \infty)$, $d \in \mathbb{N}$, let $u \in C([0, T] \times \mathbb{R}^d, \mathbb{R})$ be at most polynomially growing, assume $u|_{(0, T) \times \mathbb{R}^d} \in C^{1,2}((0, T) \times \mathbb{R}^d, \mathbb{R})$, assume for all $t \in (0, T)$, $x \in \mathbb{R}^d$ that*

$$\frac{\partial}{\partial t} u(t, x) + \frac{\kappa^2}{2} \Delta_x u(t, x) = 0, \quad (109)$$

let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $\mathbb{T}_n: \Omega \rightarrow [0, T]$, $n \in \mathbb{N}$, be *i.i.d.* random variables, let $\mathbb{X}_n: \Omega \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}$, be *i.i.d.* random variables, let $W^n: [0, T] \times \Omega \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}$, be *i.i.d.* standard Brownian motions, let $q \in [1, \infty)$ satisfy $\mathbb{E}[\|\mathbb{X}_1\|^{2q}] + \sup_{x \in \mathbb{R}^d} ((1+|x|)^{-q} |u(T, x)|) < \infty$, assume that $(\mathbb{T}_n)_{n \in \mathbb{N}}$, $(\mathbb{X}_n)_{n \in \mathbb{N}}$, and $(W^n)_{n \in \mathbb{N}}$ are independent, for every $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ let $\mathbb{F}_{M, \ell}: \mathbb{R}^{\mathfrak{d}(\ell)} \rightarrow \mathbb{R}$ satisfy for all $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ that

$$\mathbb{F}_{M, \ell}(\theta) = \frac{1}{M_1} \sum_{m=1}^{M_1} |\mathcal{N}_L^{\ell, \theta}(\mathbb{T}_m, \mathbb{X}_m) - [\frac{1}{M_2} \sum_{n=1}^{M_2} u(T, \mathbb{X}_m + \kappa W_{T-\mathbb{T}_m}^m)]|^2, \quad (110)$$

and for every $L \in \mathbb{N} \setminus \{1\}$, $\ell \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ let $\mathbf{F}_\ell: \mathbb{R}^{\mathfrak{d}(\ell)} \rightarrow [0, \infty]$ satisfy for all $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ that

$$\mathbf{F}_\ell(\theta) = \mathbb{E}[|\mathcal{N}_L^{\ell, \theta}(\mathbb{T}_1, \mathbb{X}_1) - u(\mathbb{T}_1, \mathbb{X}_1)|^2]. \quad (111)$$

Then there exists $c \in \mathbb{R}$ such that for every $R_1, R_2 \in (1, \infty)$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ and every random variable $\vartheta: \Omega \rightarrow [-R_2, R_2]^{\mathfrak{d}(\ell)}$ it holds that

$$\begin{aligned} \mathbb{E}[\mathbf{F}_\ell(\vartheta)] &\leq [\inf_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} \mathbf{F}_\ell(\theta)] + \mathbb{E}[\mathbb{F}_{M, \ell}(\vartheta) - \inf_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} \mathbb{F}_{M, \ell}(\theta)] \\ &\quad + c^L (\mathfrak{d}(\ell))^{4q} [\max\{R_1, R_2\} \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{2L} (\mathbb{E}[(1 + \|\mathbb{X}_1\|)^{2q(\mathfrak{d}(\ell))^2}])^{(\mathfrak{d}(\ell))^{-2}} (M_1)^{-1/2} \end{aligned} \quad (112)$$

(cf. Definition 1.1).

Proof of Proposition 4.5. Throughout this proof let $U: [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ satisfy for all $t \in [0, T]$, $x \in \mathbb{R}^d$ that $U(t, x) = u(t, \kappa x)$, let $G: \mathbb{R}^d \rightarrow \mathbb{R}$ satisfy for all $x \in \mathbb{R}^d$ that $G(x) = u(T, \kappa x)$, for every $n \in \mathbb{N}$ let $\mathbf{X}_n: \Omega \rightarrow \mathbb{R}^d$ satisfy $\mathbf{X}_n = \kappa^{-1} \mathbb{X}_n$, for every $L \in \mathbb{N}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in \mathbb{N}^{L+1}$, $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $\mathbb{A} \in C(\mathbb{R}, \mathbb{R})$ let $\mathbf{N}_v^{\ell, \theta, \mathbb{A}} = (\mathbf{N}_{v,1}^{\ell, \theta, \mathbb{A}}, \dots, \mathbf{N}_{v, \ell_v}^{\ell, \theta, \mathbb{A}}): \mathbb{R}^{\ell_0} \rightarrow \mathbb{R}^{\ell_v}$, $v \in \{1, 2, \dots, L\}$, satisfy for all $v \in \{0, 1, \dots, L-1\}$, $x = (x_1, \dots, x_{\ell_0}) \in \mathbb{R}^{\ell_0}$, $i \in \{1, 2, \dots, \ell_{v+1}\}$ that

$$\begin{aligned} \mathbf{N}_{v+1, i}^{\ell, \theta, \mathbb{A}}(x) &= \theta_{\ell_{v+1} \ell_v + i + \sum_{h=1}^v \ell_h (\ell_{h-1} + 1)} \\ &\quad + \sum_{j=1}^{\ell_v} \theta_{(i-1)\ell_v + j + \sum_{h=1}^v \ell_h (\ell_{h-1} + 1)} (x_j \mathbb{1}_{\{0\}}(v) + \mathbb{A}(\mathbf{N}_{\max\{v, 1\}, j}^{\ell, \theta, \mathbb{A}}(x)) \mathbb{1}_{\mathbb{N}}(v)), \end{aligned} \quad (113)$$

let $\mathcal{G} \subseteq \mathcal{F}$ satisfy

$$\mathcal{G} = \sigma((\mathbb{T}_1, \mathbf{X}_1)), \quad (114)$$

let $\mathcal{A}_n \in C^1(\mathbb{R}, \mathbb{R})$, $n \in \mathbb{N}_0$, satisfy that

- (i) it holds for all $x \in \mathbb{R}$ that $\limsup_{n \rightarrow \infty} [|\mathcal{A}_n(x) - \max\{x, 0\}| + |(\mathcal{A}_n)'(x) - \mathbb{1}_{(0, \infty)}(x)|] = 0$,
- (ii) it holds for all $x \in \mathbb{R}$ that $\mathcal{A}_0(x) = \max\{x, 0\}$, and
- (iii) it holds that $\sup_{n \in \mathbb{N}} \sup_{x \in \mathbb{R}} |(\mathcal{A}_n)'(x)| < \infty$

(cf. [66, Lemma 2.4]), for every $n \in \mathbb{N}_0$, $L \in \mathbb{N} \setminus \{1\}$, $M \in \mathbb{N}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ let $\mathbf{S}_{M, \ell}^n: \mathbb{R}^{\mathfrak{d}(\ell)} \rightarrow \mathbb{R}$ satisfy for all $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ that

$$\mathbf{S}_{M, \ell}^n(\theta) = \mathbb{E}[|\mathbf{N}_L^{\ell, \theta, \mathcal{A}_n}(\mathbb{T}_1, \kappa \mathbf{X}_1) - [\frac{1}{M} \sum_{m=1}^M G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m)]|^2], \quad (115)$$

and for every $n \in \mathbb{N}_0$, $L \in \mathbb{N} \setminus \{1\}$, $M = (M_1, M_2) \in \mathbb{N}^2$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ let $\mathbf{S}_\ell^n: \mathbb{R}^{\mathfrak{d}(\ell)} \rightarrow \mathbb{R}$ satisfy for all $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ that

$$\mathbf{S}_{M, \ell}^n(\theta) = \frac{1}{M_1} \sum_{m=1}^{M_1} |\mathbf{N}_L^{\ell, \theta, \mathcal{A}_n}(\mathbb{T}_m, \kappa \mathbf{X}_m) - [\frac{1}{M_2} \sum_{n=1}^{M_2} G(\mathbf{X}_m + W_{T-\mathbb{T}_m}^m)]|^2. \quad (116)$$

Note that Lemma 2.2, the fact that $\sup_{x \in \mathbb{R}^d} ((1 + \|x\|)^{-q} |G(x)|) < \infty$, the assumption that $\mathbb{E}[\|\mathbf{X}_1\|^{2q}] < \infty$, and the assumption that W is a standard Brownian motion demonstrate that for every $M \in \mathbb{N}$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ and every bounded set $D \subseteq \mathbb{R}^{\mathfrak{d}(\ell)}$ there exists $c \in \mathbb{R}$ such that for all $\theta \in D$ it holds that

$$\mathbb{E}[|\mathcal{N}_L^{\ell, \theta}(\mathbb{T}_1, \kappa \mathbf{X}_1)|^2] \leq c(1 + \mathbb{E}[\|\mathbb{T}_1\|^2] + \mathbb{E}[\|\mathbf{X}_1\|^2]) < \infty \quad (117)$$

and

$$\mathbb{E}\left[\left|\frac{1}{M} \sum_{m=1}^M G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m)\right|^2\right] \leq c(1 + \mathbb{E}[\|\mathbf{X}_1\|^{2q}] + \mathbb{E}[\sup_{t \in [0, T]} \|W_t\|^{2q}]) < \infty. \quad (118)$$

(cf. Definition 1.1). Combining this and (115) establishes that for all $M \in \mathbb{N}$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ it holds that

$$\begin{aligned} \mathbf{S}_{M, \ell}^0(\theta) &= \mathbb{E}\left[|\mathcal{N}_L^{\ell, \theta}(\mathbb{T}_1, \kappa \mathbf{X}_1) - \mathbb{E}\left[\frac{1}{M} \sum_{m=1}^M G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m) \middle| \mathcal{G}\right]\right. \\ &\quad \left.+ \mathbb{E}\left[\frac{1}{M} \sum_{m=1}^M G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m) \middle| \mathcal{G}\right] - \left[\frac{1}{M} \sum_{m=1}^M G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m)\right]\right|^2] \\ &= \mathbb{E}\left[|\mathcal{N}_L^{\ell, \theta}(\mathbb{T}_1, \kappa \mathbf{X}_1) - \mathbb{E}\left[\frac{1}{M} \sum_{m=1}^M G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m) \middle| \mathcal{G}\right]\right|^2] \\ &\quad + \mathbb{E}\left[\left|\mathbb{E}\left[\frac{1}{M} \sum_{m=1}^M G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m) \middle| \mathcal{G}\right] - \left[\frac{1}{M} \sum_{m=1}^M G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m)\right]\right|^2\right]. \end{aligned} \quad (119)$$

In the next step we combine (109), the assumption that for all $t \in [0, T]$, $x \in \mathbb{R}^d$ it holds that $U(t, x) = u(t, \kappa x)$, and the assumption that for all $x \in \mathbb{R}^d$ it holds that $G(x) = u(T, \kappa x)$ to obtain that for all $t \in (0, T)$, $x \in \mathbb{R}^d$ it holds that

$$\frac{\partial}{\partial t} U(t, x) + \frac{1}{2} \Delta_x U(t, x) = 0. \quad (120)$$

This, (114), the fact that $(\mathbb{T}_n)_{n \in \mathbb{N}}$, $(\mathbf{X}_n)_{n \in \mathbb{N}}$, and $(W^n)_{n \in \mathbb{N}}$ are independent, the fact that $\mathbb{E}[\|\mathbf{X}_1\|^q] < \infty$, and Lemma 4.3 ensure that for all $M \in \mathbb{N}$ it holds \mathbb{P} -a.s. that

$$\begin{aligned} \mathbb{E}\left[\frac{1}{M} \sum_{m=1}^M G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m) \middle| \mathcal{G}\right] &= \frac{1}{M} \sum_{m=1}^M \mathbb{E}[G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m) | (\mathbb{T}_1, \mathbf{X}_1)] \\ &= \frac{1}{M} \sum_{m=1}^M U(\mathbb{T}_1, \mathbf{X}_1) = U(\mathbb{T}_1, \mathbf{X}_1). \end{aligned} \quad (121)$$

Combining this and (119) implies for all $M \in \mathbb{N}$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ that

$$\begin{aligned} \mathbf{S}_{M, \ell}^0(\theta) &= \mathbb{E}[|\mathcal{N}_L^{\ell, \theta}(\mathbb{T}_1, \kappa \mathbf{X}_1) - U(\mathbb{T}_1, \mathbf{X}_1)|^2] \\ &\quad + \mathbb{E}\left[\left|U(\mathbb{T}_1, \mathbf{X}_1) - \left[\frac{1}{M} \sum_{m=1}^M G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m)\right]\right|^2\right]. \end{aligned} \quad (122)$$

Next, observe that (111), the fact that $\mathbb{X}_1 = \kappa \mathbf{X}_1$, and the assumption that for all $t \in [0, T]$, $x \in \mathbb{R}^d$ it holds that $U(t, x) = u(t, \kappa x)$ show that for all $L \in \mathbb{N} \setminus \{1\}$, $\ell \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ it holds that

$$\mathbf{F}_\ell(\theta) = \mathbb{E}[|\mathcal{N}_L^{\ell, \theta}(\mathbb{T}_1, \kappa \mathbf{X}_1) - U(\mathbb{T}_1, \mathbf{X}_1)|^2]. \quad (123)$$

This and (122) prove that for all $M \in \mathbb{N}$, $R \in (0, \infty]$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ it holds that

$$\begin{aligned} \inf_{\theta \in (-R, R)^{\mathfrak{d}(\ell)}} \mathbf{S}_{M, \ell}^0(\theta) &= \inf_{\theta \in (-R, R)^{\mathfrak{d}(\ell)}} \mathbf{F}_\ell(\theta) \\ &\quad + \mathbb{E}\left[\left|U(\mathbb{T}_1, \mathbf{X}_1) - \left[\frac{1}{M} \sum_{m=1}^M G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m)\right]\right|^2\right]. \end{aligned} \quad (124)$$

Hence, we obtain for all $M \in \mathbb{N}$, $R \in (0, \infty]$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ that

$$\begin{aligned} &\mathbb{E}\left[\left|U(\mathbb{T}_1, \mathbf{X}_1) - \left[\frac{1}{M} \sum_{m=1}^M G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m)\right]\right|^2\right] \\ &= \inf_{\theta \in (-R, R)^{\mathfrak{d}(\ell)}} \mathbf{S}_{M, \ell}^0(\theta) - \inf_{\theta \in (-R, R)^{\mathfrak{d}(\ell)}} \mathbf{F}_\ell(\theta). \end{aligned} \quad (125)$$

Combining this and (122) demonstrates for all $M \in \mathbb{N}$, $R \in (0, \infty]$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $\vartheta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ that

$$\begin{aligned} \mathbf{S}_{M,\ell}^0(\vartheta) &= \mathbb{E}[|\mathcal{N}_L^{\ell,\vartheta}(\mathbb{T}_1, \kappa \mathbf{X}_1) - U(\mathbb{T}_1, \mathbf{X}_1)|^2] \\ &\quad + [\inf_{\theta \in (-R,R)^{\mathfrak{d}(\ell)}} \mathbf{S}_{M,\ell}^0(\theta)] - [\inf_{\theta \in (-R,R)^{\mathfrak{d}(\ell)}} \mathbf{F}_\ell(\theta)]. \end{aligned} \quad (126)$$

This establishes for all $M \in \mathbb{N}$, $R \in (0, \infty]$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $\vartheta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ that

$$\begin{aligned} \mathbb{E}[|\mathcal{N}_L^{\ell,\vartheta}(\mathbb{T}_1, \kappa \mathbf{X}_1) - U(\mathbb{T}_1, \mathbf{X}_1)|^2] &\leq \mathbf{S}_{M,\ell}^0(\vartheta) - \inf_{\theta \in (-R,R)^{\mathfrak{d}(\ell)}} \mathbf{S}_{M,\ell}^0(\theta) \\ &\quad + \inf_{\theta \in (-R,R)^{\mathfrak{d}(\ell)}} \mathbf{F}_\ell(\theta). \end{aligned} \quad (127)$$

Combining this and (111) ensures for all $M \in \mathbb{N}^2$, $R \in (0, \infty]$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $\vartheta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ that

$$\mathbf{F}_\ell(\vartheta) \leq [\inf_{\theta \in (-R,R)^{\mathfrak{d}(\ell)}} \mathbf{F}_\ell(\theta)] + (\mathbf{S}_{M,\ell}^0(\vartheta) - \inf_{\theta \in (-R,R)^{\mathfrak{d}(\ell)}} \mathbf{S}_{M,\ell}^0(\theta)). \quad (128)$$

The fact that for all $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ it holds that $\mathbf{S}_{M_2,\ell}^0$ and $\mathbb{F}_{M,\ell}$ are continuous hence implies that for all $M \in \mathbb{N}^2$, $R_1 \in (1, \infty]$, $R_2 \in (1, \infty)$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $\vartheta \in [-R_2, R_2]^{\mathfrak{d}(\ell)}$ it holds that

$$\begin{aligned} \mathbf{F}_\ell(\vartheta) &\leq [\inf_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} \mathbf{F}_\ell(\theta)] + (\mathbb{F}_{M,\ell}(\vartheta) - \inf_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} \mathbb{F}_{M,\ell}(\theta)) \\ &\quad + [\sup_{\theta \in (-R_2, R_2)^{\mathfrak{d}(\ell)}} |\mathbf{S}_{M_2,\ell}^0(\theta) - \mathbb{F}_{M,\ell}(\theta)|] + [\sup_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} |\mathbf{S}_{M_2,\ell}^0(\theta) - \mathbb{F}_{M,\ell}(\theta)|]. \end{aligned} \quad (129)$$

In addition, note that (110), (117), (118), the assumption that $\sup_{x \in \mathbb{R}^d} ((1+|x|)^{-q} |u(T, x)|) < \infty$, the assumption that W is a standard Brownian motion, and the assumption that $\mathbb{P}(\mathbb{T} \leq T) = 1$ show that for all $R \in (0, \infty]$, $M \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ there exists $c \in (0, \infty)$ such that

$$\begin{aligned} &\mathbb{E}[\inf_{\theta \in (-R,R)^{\mathfrak{d}(\ell)}} \mathbb{F}_{M,\ell}(\theta)] \\ &\leq \mathbb{E}[\mathbb{F}_{M,\ell}(0)] \leq c(1 + \mathbb{E}[\|\mathbf{X}_1\|^{2q}] + \mathbb{E}[\sup_{t \in [0,T]} \|W_t\|^{2q}]) < \infty. \end{aligned} \quad (130)$$

This and (129) prove that for every $M \in \mathbb{N}^2$, $R_1 \in (1, \infty]$, $R_2 \in (1, \infty)$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ and every random variable $\vartheta: \Omega \rightarrow [-R_2, R_2]^{\mathfrak{d}(\ell)}$ it holds that

$$\begin{aligned} \mathbb{E}[\mathbf{F}_\ell(\vartheta)] &\leq \mathbb{E}[\inf_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} \mathbf{F}_\ell(\theta)] + \mathbb{E}[\mathbb{F}_{M,\ell}(\vartheta) - \inf_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} \mathbb{F}_{M,\ell}(\theta)] \\ &\quad + \mathbb{E}[\sup_{\theta \in (-R_2, R_2)^{\mathfrak{d}(\ell)}} |\mathbf{S}_{M_2,\ell}^0(\theta) - \mathbb{F}_{M,\ell}(\theta)|] + \mathbb{E}[\sup_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} |\mathbf{S}_{M_2,\ell}^0(\theta) - \mathbb{F}_{M,\ell}(\theta)|]. \end{aligned} \quad (131)$$

Moreover, observe that (115), (116), and Corollary 4.2 demonstrate that there exists $\mathfrak{C} \in \mathbb{R}$ which satisfies for all $n \in \mathbb{N}$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $\vartheta \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $p \in (\mathfrak{d}(\ell))^2, \infty)$, $R \in [1/2, \infty)$ that

$$\begin{aligned} &\mathbb{E}[\sup_{\theta \in (-R,R)^{\mathfrak{d}(\ell)}} |\mathbf{S}_{M_2,\ell}^n(\theta) - \mathbb{S}_{M,\ell}^n(\theta)|^p] \\ &= \mathbb{E}[\sup_{\theta \in (-R,R)^{\mathfrak{d}(\ell)}} |\mathbb{S}_{M,\ell}^n(\theta) - \mathbb{E}[\mathbb{S}_{M,\ell}^n(\theta)]|^p] \\ &\leq |\mathfrak{R}\mathfrak{C}|^p \mathbb{E} \left[\int_{[-R,R]^{\mathfrak{d}(\ell)}} (|\mathbb{S}_{M,\ell}^n(\theta) - \mathbb{E}[\mathbb{S}_{M,\ell}^n(\theta)]|^p + \|(\nabla \mathbb{S}_{M,\ell}^n)(\theta) - \mathbb{E}[(\nabla \mathbb{S}_{M,\ell}^n)(\theta)]\|^p) d\theta \right] \\ &= |\mathfrak{R}\mathfrak{C}|^p \int_{[-R,R]^{\mathfrak{d}(\ell)}} (\mathbb{E}[|\mathbb{S}_{M,\ell}^n(\theta) - \mathbb{E}[\mathbb{S}_{M,\ell}^n(\theta)]|^p] + \mathbb{E}[\|(\nabla \mathbb{S}_{M,\ell}^n)(\theta) - \mathbb{E}[(\nabla \mathbb{S}_{M,\ell}^n)(\theta)]\|^p]) d\theta. \end{aligned} \quad (132)$$

Note that (132) establishes that for all $n \in \mathbb{N}$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $p \in (\mathfrak{d}(\ell))^2, \infty)$, $R \in [1/2, \infty)$ it holds that

$$\begin{aligned} & \mathbb{E} \left[\sup_{\theta \in (-R, R)^{\mathfrak{d}(\ell)}} |\mathbf{S}_{M_2, \ell}^n(\theta) - \mathbb{S}_{M, \ell}^n(\theta)|^p \right] \\ & \leq |R\mathfrak{C}|^p (2R)^{\mathfrak{d}(\ell)} \sup_{\theta \in (-R, R)^{\mathfrak{d}(\ell)}} \mathbb{E} \left[|\mathbb{S}_{M, \ell}^n(\theta) - \mathbb{E}[\mathbb{S}_{M, \ell}^n(\theta)]|^p \right] \\ & \quad + |R\mathfrak{C}|^p (2R)^{\mathfrak{d}(\ell)} \sup_{\theta \in (-R, R)^{\mathfrak{d}(\ell)}} \mathbb{E} \left[\left\| (\nabla \mathbb{S}_{M, \ell}^n)(\theta) - \mathbb{E}[(\nabla \mathbb{S}_{M, \ell}^n)(\theta)] \right\|^p \right]. \end{aligned} \quad (133)$$

Furthermore, observe that Lemma 4.4, (116), the Cauchy–Schwarz inequality, the fact that for all $x, y \in \mathbb{R}^d$, $p \in [1, \infty)$ it holds that $\|x+y\|^p \leq 2^{p-1}\|x\|^p + 2^{p-1}\|y\|^p$, and the fact that for every $p \in [2, \infty)$ and every random variable $X: \Omega \rightarrow \mathbb{R}^{\mathfrak{d}}$ it holds that $\mathbb{E}[\|X - \mathbb{E}[X]\|^p] \leq 2^p \mathbb{E}[\|X\|^p]$ ensure that for all $n \in \mathbb{N}_0$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $\vartheta \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $p \in [2, \infty)$ it holds that

$$\begin{aligned} & \mathbb{E} \left[|\mathbb{S}_{M, \ell}^n(\vartheta) - \mathbb{E}[\mathbb{S}_{M, \ell}^n(\vartheta)]|^p \right] \\ & \leq 2^p \left[\frac{p-1}{M_1} \right]^{p/2} \mathbb{E} \left[\left| (\mathbf{N}_L^{\ell, \vartheta, \mathcal{A}_n}(\mathbb{T}_1, \kappa \mathbf{X}_1) - \left[\frac{1}{M_2} \sum_{m=1}^{M_2} G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m) \right]) \right. \right. \\ & \quad \left. \left. - \mathbb{E} \left[(\mathbf{N}_L^{\ell, \vartheta, \mathcal{A}_n}(\mathbb{T}_1, \kappa \mathbf{X}_1) - \left[\frac{1}{M_2} \sum_{m=1}^{M_2} G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m) \right]) \right] \right|^p \right] \\ & \leq 2^{2p} \left[\frac{p-1}{M_1} \right]^{p/2} \mathbb{E} \left[\left| \mathbf{N}_L^{\ell, \vartheta, \mathcal{A}_n}(\mathbb{T}_1, \kappa \mathbf{X}_1) - \left[\frac{1}{M_2} \sum_{m=1}^{M_2} G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m) \right] \right|^{2p} \right] \\ & \leq 2^{4p-1} \left[\frac{p-1}{M_1} \right]^{p/2} \left(\mathbb{E} \left[|\mathbf{N}_L^{\ell, \vartheta, \mathcal{A}_n}(\mathbb{T}_1, \kappa \mathbf{X}_1)|^{2p} \right] + \mathbb{E} \left[\left| \frac{1}{M_2} \sum_{m=1}^{M_2} G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^m) \right|^{2p} \right] \right). \end{aligned} \quad (134)$$

In the next step we note that (116) implies that for all $n \in \mathbb{N}$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $N = (N_1, N_2) \in \mathbb{N}^2$, $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ it holds that

$$\begin{aligned} (\nabla \mathbb{S}_{N, \ell}^n)(\theta) &= \frac{1}{N_1} \left(\sum_{m=1}^{N_1} 2(\mathbf{N}_L^{\ell, \theta, \mathcal{A}_n}(\mathbb{T}_m, \kappa \mathbf{X}_m) - \left[\frac{1}{N_2} \sum_{n=1}^{N_2} G(\mathbf{X}_m + W_{T-\mathbb{T}_m}^{N_2+n}) \right]) \right. \\ & \quad \left. \cdot [\nabla_{\theta}(\mathbf{N}_L^{\ell, \theta, \mathcal{A}_n}(\mathbb{T}_m, \kappa \mathbf{X}_m))] \right). \end{aligned} \quad (135)$$

Combining this, Lemma 4.4, the Cauchy–Schwarz inequality, the fact that for all $x, y \in \mathbb{R}^d$, $p \in [1, \infty)$ it holds that $\|x+y\|^p \leq 2^{p-1}\|x\|^p + 2^{p-1}\|y\|^p$, and the fact that for every $p \in [2, \infty)$ and every random variable $X: \Omega \rightarrow \mathbb{R}^{\mathfrak{d}}$ it holds that $\mathbb{E}[\|X - \mathbb{E}[X]\|^p] \leq 2^p \mathbb{E}[\|X\|^p]$ shows that for all $n \in \mathbb{N}$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $\vartheta \in \mathbb{R}^{\mathfrak{d}(\ell)}$, $p \in [2, \infty)$ it holds that

$$\begin{aligned} & \mathbb{E} \left[\left\| (\nabla \mathbb{S}_{M, \ell}^n)(\vartheta) - \mathbb{E}[(\nabla \mathbb{S}_{M, \ell}^n)(\vartheta)] \right\|^p \right] \\ & \leq 2^p \left[\frac{p-1}{M_1} \right]^{p/2} \mathbb{E} \left[\left\| 2(\mathbf{N}_L^{\ell, \vartheta, \mathcal{A}_n}(\mathbb{T}_1, \kappa \mathbf{X}_1) - \left[\frac{1}{M_2} \sum_{n=1}^{M_2} G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^n) \right]) \right. \right. \\ & \quad \left. \left. - \mathbb{E} \left[2(\mathbf{N}_L^{\ell, \vartheta, \mathcal{A}_n}(\mathbb{T}_1, \kappa \mathbf{X}_1) - \left[\frac{1}{M_2} \sum_{n=1}^{M_2} G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^n) \right]) \right] \right\|^p \right] \\ & \leq 2^{3p} \left[\frac{p-1}{M_1} \right]^{p/2} \mathbb{E} \left[\left\| (\mathbf{N}_L^{\ell, \vartheta, \mathcal{A}_n}(\mathbb{T}_1, \kappa \mathbf{X}_1) - \left[\frac{1}{M_2} \sum_{n=1}^{M_2} G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^n) \right]) \right\|^p \right] \\ & \leq 2^{3p+1} \left[\frac{p-1}{M_1} \right]^{p/2} \mathbb{E} \left[\left| \mathbf{N}_L^{\ell, \vartheta, \mathcal{A}_n}(\mathbb{T}_1, \kappa \mathbf{X}_1) - \left[\frac{1}{M_2} \sum_{n=1}^{M_2} G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^n) \right] \right|^{2p} \right] \\ & \quad + 2^{3p+1} \left[\frac{p-1}{M_1} \right]^{p/2} \mathbb{E} \left[\left\| \nabla_{\vartheta}(\mathbf{N}_L^{\ell, \vartheta, \mathcal{A}_n}(\mathbb{T}_1, \kappa \mathbf{X}_1)) \right\|^{2p} \right] \\ & \leq 2^{5p} \left[\frac{p-1}{M_1} \right]^{p/2} \mathbb{E} \left[|\mathbf{N}_L^{\ell, \vartheta, \mathcal{A}_n}(\mathbb{T}_1, \kappa \mathbf{X}_1)|^{2p} \right] + 2^{5p} \left[\frac{p-1}{M_1} \right]^{p/2} \mathbb{E} \left[\left| \frac{1}{M_2} \sum_{n=1}^{M_2} G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^n) \right|^{2p} \right] \\ & \quad + 2^{3p+1} \left[\frac{p-1}{M_1} \right]^{p/2} \mathbb{E} \left[\left\| \nabla_{\vartheta}(\mathbf{N}_L^{\ell, \vartheta, \mathcal{A}_n}(\mathbb{T}_1, \kappa \mathbf{X}_1)) \right\|^{2p} \right]. \end{aligned} \quad (136)$$

Next we combine the fact that $\sup_{x \in \mathbb{R}^d} (1+\|x\|)^{-q} |G(x)| < \infty$, the assumption that W^m , $m \in \mathbb{N}$, are i.i.d. standard Brownian motions, Doob’s inequality, and Cauchy–Schwarz inequality to

obtain that there exist $c_1, c_2, c_3 \in \mathbb{R}$ such that for all $M \in \mathbb{N}$, $p \in [2, \infty)$ it holds that

$$\begin{aligned}
& \mathbb{E}\left[\left|\frac{1}{M} \sum_{n=1}^M G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^n)\right|^{2p}\right] \\
& \leq \frac{1}{M} \sum_{n=1}^M \mathbb{E}\left[|G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^n)|^{2p}\right] = \mathbb{E}\left[|G(\mathbf{X}_1 + W_{T-\mathbb{T}_1}^1)|^{2p}\right] \\
& \leq |c_1|^{2pq} + |c_1|^{2pq} \mathbb{E}\left[\|\mathbf{X}_1\|^{2pq}\right] + |c_1|^{2pq} \mathbb{E}\left[\sup_{t \in [0, T]} |W_t^1|^{2pq}\right] \\
& \leq |c_2|^{2pq} + |c_2|^{2pq} \mathbb{E}\left[\|\mathbf{X}_1\|^{2pq}\right] + |c_2|^{2pq} \left(\frac{2pq}{2pq-1}\right)^{2p} T^{2p} (\lceil 2pq \rceil)!! \\
& \leq |2c_3|^{2pq} (\mathbb{E}\left[\|\mathbf{X}_1\|^{2pq}\right] + (2pq)^{pq}) \leq |c_3|^{2pq} (pq)^{pq} \mathbb{E}\left[(1 + \|\mathbf{X}_1\|)^{2pq}\right].
\end{aligned} \tag{137}$$

Next, observe that Corollary 2.4 and item (iii) prove that there exist $c_1, c_2 \in (0, \infty)$ such that for all $R \in (1, \infty]$, $n \in \mathbb{N}$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $\vartheta = (\vartheta_1, \dots, \vartheta_{\mathfrak{d}(\ell)}) \in [-R, R]^{\mathfrak{d}(\ell)}$, $p \in [2, \infty)$ it holds that

$$\begin{aligned}
& \mathbb{E}\left[\left\|\nabla_{\vartheta}(\mathbf{N}_L^{\ell, \vartheta, \mathcal{A}_n}(\mathbb{T}_1, \kappa \mathbf{X}_1))\right\|^{2p}\right] \\
& \leq \mathbb{E}\left[\max\{1, \|(\mathbb{T}_1, \kappa \mathbf{X}_1)\|^{2p}\} \max\left\{1, \max_{j \in \{1, 2, \dots, \mathfrak{d}(\ell)\}} |\vartheta_j|^{2pL}\right\} \left[3c_1 \max\{\ell_0, \ell_1, \dots, \ell_{L-1}\}\right]^{2pL}\right] \\
& \leq (c_2)^{2pL} R^{2pL} [\max\{\ell_0, \ell_1, \dots, \ell_{L-1}\}]^{2pL} \mathbb{E}\left[(1 + \|\mathbf{X}_1\|)^{2p}\right] \\
& = (c_2)^{2pL} [R \max\{\ell_0, \ell_1, \dots, \ell_{L-1}\}]^{2pL} \mathbb{E}\left[(1 + \|\mathbf{X}_1\|)^{2p}\right].
\end{aligned} \tag{138}$$

In addition, note that Lemma 2.2 and item (iii) demonstrate that there exist $c_1, c_2 \in (0, \infty)$ such that for all $R \in (1, \infty]$, $n \in \mathbb{N}$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $\vartheta = (\vartheta_1, \dots, \vartheta_{\mathfrak{d}(\ell)}) \in [-R, R]^{\mathfrak{d}(\ell)}$, $p \in [2, \infty)$ it holds that

$$\begin{aligned}
& \mathbb{E}\left[\left|\mathbf{N}_L^{\ell, \vartheta, \mathcal{A}_n}(\mathbb{T}_1, \kappa \mathbf{X}_1)\right|^{2p}\right] \\
& \leq \mathbb{E}\left[\max\{1, \|(\mathbb{T}_1, \kappa \mathbf{X}_1)\|^{2p}\} \max\left\{1, \max_{j \in \{1, 2, \dots, \mathfrak{d}(\ell)\}} |\vartheta_j|^{2pL}\right\} \left[3c_1 \max\{\ell_0, \ell_1, \dots, \ell_{L-1}\}\right]^{2pL}\right] \\
& \leq (c_2)^{2pL} R^{2pL} [\max\{\ell_0, \ell_1, \dots, \ell_{L-1}\}]^{2pL} \mathbb{E}\left[(1 + \|\mathbf{X}_1\|)^{2p}\right] \\
& \leq (c_3)^{2pL} [R \max\{\ell_0, \ell_1, \dots, \ell_{L-1}\}]^{2pL} \mathbb{E}\left[(1 + \|\mathbf{X}_1\|)^{2p}\right].
\end{aligned} \tag{139}$$

This, (134), (136), (137), (138), and the fact that $q \geq 1$ establish that there exists $c \in (0, \infty)$ such that for all $R \in (1, \infty]$, $n \in \mathbb{N}$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $\vartheta = (\vartheta_1, \dots, \vartheta_{\mathfrak{d}(\ell)}) \in [-R, R]^{\mathfrak{d}(\ell)}$, $p \in [2, \infty)$ it holds that

$$\begin{aligned}
& \sup_{\vartheta \in [-R, R]^{\mathfrak{d}(\ell)}} \mathbb{E}\left[|\mathbb{S}_{M, \ell}^n(\vartheta) - \mathbb{E}[\mathbb{S}_{M, \ell}^n(\vartheta)]|^p\right] + \sup_{\vartheta \in [-R, R]^{\mathfrak{d}(\ell)}} \mathbb{E}\left[\|(\nabla \mathbb{S}_{M, \ell}^n)(\vartheta) - \mathbb{E}[(\nabla \mathbb{S}_{M, \ell}^n)(\vartheta)]\|^p\right] \\
& \leq c^{2pL} (pq)^{pq} [R \max\{\ell_0, \ell_1, \dots, \ell_{L-1}\}]^{2pL} \mathbb{E}\left[(1 + \|\mathbf{X}_1\|)^{2pq}\right] (M_1)^{-p/2}.
\end{aligned} \tag{140}$$

Combining this and (133) ensures that there exists $c \in \mathbb{R}$ such that for all $n \in \mathbb{N}$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $R \in (1, \infty]$, $p \in [(\mathfrak{d}(\ell))^2, \infty)$ it holds that

$$\begin{aligned}
& \mathbb{E}\left[\sup_{\theta \in (-R, R)^{\mathfrak{d}(\ell)}} |\mathbf{S}_{M_2, \ell}^n(\theta) - \mathbb{S}_{M, \ell}^n(\theta)|^p\right] \\
& \leq c^{2pLc} (p^2 q)^{pq} [R \max\{\ell_0, \ell_1, \dots, \ell_{L-1}\}]^{2pL} \mathbb{E}\left[(1 + \|\mathbf{X}_1\|)^{2pq}\right] (M_1)^{-p/2}.
\end{aligned} \tag{141}$$

This, item (i), item (ii), item (iii), and the fact that for all $M \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell \in (\{d\} \times \mathbb{N}^{L-1} \times \{1\})$, $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ it holds that $\mathbb{F}_{M, \ell}(\theta) = \mathbb{S}_{M, \ell}^0(\theta)$ imply that there exists $c \in \mathbb{R}$ such that for all $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $R \in (1, \infty]$, $p \in [(\mathfrak{d}(\ell))^2, \infty)$ it holds that

$$\begin{aligned}
& \mathbb{E}\left[\sup_{\theta \in (-R, R)^{\mathfrak{d}(\ell)}} |\mathbf{S}_{M_2, \ell}^0(\theta) - \mathbb{F}_{M, \ell}(\theta)|^p\right] \\
& \leq \mathbb{E}\left[\sup_{\theta \in (-R, R)^{\mathfrak{d}(\ell)}} |\mathbf{S}_{M_2, \ell}^0(\theta) - \mathbb{S}_{M, \ell}^0(\theta)|^p\right] \\
& \leq c^{2pLc} (p^2 q)^{pq} [R \max\{\ell_0, \ell_1, \dots, \ell_{L-1}\}]^{2pL} \mathbb{E}\left[(1 + \|\mathbf{X}_1\|)^{2pq}\right] (M_1)^{-p/2}.
\end{aligned} \tag{142}$$

Holder's inequality therefore shows that there exists $c \in \mathbb{R}$ such that for all $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $R \in (1, \infty]$ it holds that

$$\begin{aligned} & \mathbb{E}[\sup_{\theta \in (-R, R)^{\mathfrak{d}(\ell)}} |\mathbf{S}_{M_2, \ell}^0(\theta) - \mathbb{F}_{M, \ell}(\theta)|] \\ & \leq c^{2L} (\mathfrak{d}(\ell))^{4q} [R \max\{\ell_0, \ell_1, \dots, \ell_{L-1}\}]^{2L} (\mathbb{E}[(1 + \|\mathbf{X}_1\|)^{2q(\mathfrak{d}(\ell))^2}])^{(\mathfrak{d}(\ell))^{-2}} (M_1)^{-1/2}. \end{aligned} \quad (143)$$

Combining this and the fact that for all $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ it holds that $\max\{\ell_0, \ell_1, \dots, \ell_{L-1}\} \leq (d+1) \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}$ proves that there exists $c \in \mathbb{R}$ which satisfies for all $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $R \in (1, \infty]$ that

$$\begin{aligned} & \mathbb{E}[\sup_{\theta \in (-R, R)^{\mathfrak{d}(\ell)}} |\mathbf{S}_{M_2, \ell}^0(\theta) - \mathbb{F}_{M, \ell}(\theta)|] \\ & \leq c^{2L} (\mathfrak{d}(\ell))^{4q} [R \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{2L} (\mathbb{E}[(1 + \|\mathbf{X}_1\|)^{2q(\mathfrak{d}(\ell))^2}])^{(\mathfrak{d}(\ell))^{-2}} (M_1)^{-1/2}. \end{aligned} \quad (144)$$

This and (131) demonstrate (112). The proof of Proposition 4.5 is thus complete. \square

Theorem 4.6. *Let $T, \kappa \in (0, \infty)$, $d \in \mathbb{N}$, let $u \in C^2([0, T] \times \mathbb{R}^d, \mathbb{R})$ be at most polynomially growing, assume for all $t \in (0, T)$, $x \in \mathbb{R}^d$ that*

$$\frac{\partial}{\partial t} u(t, x) + \frac{\kappa^2}{2} \Delta_x u(t, x) = 0, \quad (145)$$

let $q \in [1, \infty)$ satisfy $\sup_{x \in \mathbb{R}^d} ((1 + |x|)^{-q} |u(T, x)|) < \infty$, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $\mathbb{T}_n: \Omega \rightarrow [0, T]$, $n \in \mathbb{N}$, be *i.i.d.* random variables, let $\mathbb{X}_n: \Omega \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}$, be *i.i.d.* random variables, let $W^n: [0, T] \times \Omega \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}$, be *i.i.d.* standard Brownian motions, let $D \subseteq (0, T) \times \mathbb{R}^d$ be non-empty, open, and bounded, assume that $(\mathbb{T}_1, \mathbb{X}_1)$ is uniformly distributed on D , assume that $(\mathbb{T}_n)_{n \in \mathbb{N}}$, $(\mathbb{X}_n)_{n \in \mathbb{N}}$, and $(W^n)_{n \in \mathbb{N}}$ are independent, and for every $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ let $\mathbb{F}_{M, \ell}: \mathbb{R}^{\mathfrak{d}(\ell)} \rightarrow \mathbb{R}$ satisfy for all $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ that

$$\mathbb{F}_{M, \ell}(\theta) = \frac{1}{M_1} \sum_{m=1}^{M_1} |\mathcal{N}_L^{\ell, \theta}(\mathbb{T}_m, \mathbb{X}_m) - [\frac{1}{M_2} \sum_{n=1}^{M_2} u(T, \mathbb{X}_m + \kappa W_{T-\mathbb{T}_m}^{m M_2 + n})]|^2 \quad (146)$$

(cf. Definition 1.1). Then there exists $c \in \mathbb{R}$ such that for every $R_1, R_2 \in (1, \infty)$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ and every random variable $\vartheta: \Omega \rightarrow [-R_2, R_2]^{\mathfrak{d}(\ell)}$ it holds that

$$\begin{aligned} & \mathbb{E}[\int_D |u(y) - \mathcal{N}_L^{\ell, \vartheta}(y)|^2 dy] \leq c \mathbb{1}_{(R_1, \infty)}(c) + \lambda(D) \mathbb{E}[\mathbb{F}_{M, \ell}(\vartheta) - \inf_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} \mathbb{F}_{M, \ell}(\theta)] \\ & + c^L (\mathfrak{d}(\ell))^{4q} [\max\{R_1, R_2\} \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{2L} (M_1)^{-1/2} + c [\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{-\frac{2}{d+5}} \end{aligned} \quad (147)$$

(cf. Definition 1.1).

Proof of Theorem 4.6. Throughout this proof for every $L \in \mathbb{N} \setminus \{1\}$, $\ell \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ let $\mathbf{F}_\ell: \mathbb{R}^{\mathfrak{d}(\ell)} \rightarrow [0, \infty]$ satisfy for all $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ that

$$\mathbf{F}_\ell(\theta) = \mathbb{E}[|\mathcal{N}_L^{\ell, \theta}(\mathbb{T}_1, \mathbb{X}_1) - u(\mathbb{T}_1, \mathbb{X}_1)|^2]. \quad (148)$$

Observe that (148), Proposition 4.5, and the fact that \mathbb{X}_1 is a bounded random variable establish there exists $c \in (0, \infty)$ such that for every $R_1, R_2 \in (1, \infty)$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ and every random variable $\vartheta: \Omega \rightarrow [-R_2, R_2]^{\mathfrak{d}(\ell)}$ it holds that

$$\begin{aligned} & \mathbb{E}[\mathbf{F}_\ell(\vartheta)] \leq [\inf_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} \mathbf{F}_\ell(\theta)] + \mathbb{E}[\mathbb{F}_{M, \ell}(\vartheta) - \inf_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} \mathbb{F}_{M, \ell}(\theta)] \\ & + c^L (\mathfrak{d}(\ell))^{4q} [\max\{R_1, R_2\} \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{2L} (M_1)^{-1/2}. \end{aligned} \quad (149)$$

(cf. Definition 1.1). Moreover, note that (148) and the assumption that (\mathbb{T}, \mathbb{X}) is uniformly distributed on D ensure that for all $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$, $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ it holds that

$$\mathbf{F}_\ell(\theta) = \frac{1}{\lambda(D)} \int_D |u(y) - \mathcal{N}_L^{\ell, \theta}(y)|^2 dy. \quad (150)$$

Combining this and (149) implies that there exists $c \in (0, \infty)$ such that for every $R_1, R_2 \in (1, \infty)$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ and every random variable $\vartheta: \Omega \rightarrow [-R_2, R_2]^{\mathfrak{d}(\ell)}$ it holds that

$$\begin{aligned} & \mathbb{E} \left[\int_D |u(y) - \mathcal{N}_L^{\ell, \vartheta}(y)|^2 dy \right] \\ & \leq \left[\inf_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} (\lambda(D) \mathbf{F}_\ell(\theta)) \right] + \lambda(D) \mathbb{E} \left[\mathbb{F}_{M, \ell}(\vartheta) - \inf_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} \mathbb{F}_{M, \ell}(\theta) \right] \\ & + (\lambda(D)) c^L (\mathfrak{d}(\ell))^{4q} \left[\max\{R_1, R_2\} \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\} \right]^{2L} (M_1)^{-1/2}. \end{aligned} \quad (151)$$

Furthermore, observe that the assumption that D is open and bounded and Corollary 3.7 (applied with $d \curvearrowright d$, $D \curvearrowright D$, $u \curvearrowright u|_{\bar{D}}$ in the notation of Corollary 3.7) show that there exists $c \in (0, \infty)$ which satisfies for all $R \in (0, \infty]$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ that

$$\inf_{\theta \in (-R, R)^{\mathfrak{d}(\ell)}} \int_D |u(y) - \mathcal{N}_L^{\ell, \theta}(y)|^2 \leq c \mathbb{1}_{(-\infty, c)}(R) + c [\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{\frac{-2}{d+5}}. \quad (152)$$

Note that (150) and (152) prove that for all $R \in (0, \infty]$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ it holds that

$$\inf_{\theta \in (-R, R)^{\mathfrak{d}(\ell)}} (\lambda(D) \mathbf{F}_\ell(\theta)) \leq \mathbb{1}_{(R, \infty)}(c) + [\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{\frac{-2}{d+5}}. \quad (153)$$

This and (151) demonstrate (147). The proof of Theorem 4.6 is thus complete. \square

Corollary 4.7. *Let $T, \kappa \in (0, \infty)$, $d \in \mathbb{N}$, let $u \in C^2([0, T] \times \mathbb{R}^d, \mathbb{R})$ be at most polynomially growing, assume for all $t \in (0, T)$, $x \in \mathbb{R}^d$ that*

$$\frac{\partial}{\partial t} u(t, x) + \frac{\kappa^2}{2} \Delta_x u(t, x) = 0, \quad (154)$$

let $q \in [1, \infty)$ satisfy $\sup_{x \in \mathbb{R}^d} ((1 + |x|)^{-q} |u(T, x)|) < \infty$, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $\mathbb{T}_n: \Omega \rightarrow [0, T]$, $n \in \mathbb{N}$, be *i.i.d.* random variables, let $\mathbb{X}_n: \Omega \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}$, be *i.i.d.* random variables, let $W^n: [0, T] \times \Omega \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}$, be *i.i.d.* standard Brownian motions, let $D \subseteq (0, T) \times \mathbb{R}^d$ be non-empty, open, and bounded, assume that $(\mathbb{T}_1, \mathbb{X}_1)$ is uniformly distributed on D , assume that $(\mathbb{T}_n)_{n \in \mathbb{N}}$, $(\mathbb{X}_n)_{n \in \mathbb{N}}$, and $(W^n)_{n \in \mathbb{N}}$ are independent, and for every $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ let $\mathbb{F}_{M, \ell}: \mathbb{R}^{\mathfrak{d}(\ell)} \rightarrow \mathbb{R}$ satisfy for all $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ that

$$\mathbb{F}_{M, \ell}(\theta) = \frac{1}{M_1} \sum_{m=1}^{M_1} |\mathcal{N}_L^{\ell, \theta}(\mathbb{T}_m, \mathbb{X}_m) - \left[\frac{1}{M_2} \sum_{n=1}^{M_2} u(T, \mathbb{X}_m + \kappa W_{T-\mathbb{T}_m}^{m, M_2+n}) \right]|^2 \quad (155)$$

(cf. Definition 1.1). Then there exists $c \in \mathbb{R}$ such that for every $R_1, R_2 \in (1, \infty)$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ and every random variable $\vartheta: \Omega \rightarrow [-R_2, R_2]^{\mathfrak{d}(\ell)}$ it holds that

$$\begin{aligned} & \mathbb{E} \left[\int_D |u(y) - \mathcal{N}_L^{\ell, \vartheta}(y)|^2 dy \right] \\ & \leq c^L (\mathfrak{d}(\ell))^{4q} \left[\max\{R_1, R_2\} \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\} \right]^{2L} (M_1)^{-1/2} + c \exp(-R_1) \mathbb{1}_{(-\infty, c)}(R_1) \\ & + c [\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{\frac{-2}{d+5}} + \lambda(D) \mathbb{E} \left[\mathbb{F}_{M, \ell}(\vartheta) - \inf_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} \mathbb{F}_{M, \ell}(\theta) \right] \end{aligned} \quad (156)$$

(cf. Definition 1.1).

Proof of Corollary 4.7. Observe that Theorem 4.6 establishes that there exists $c \in (0, \infty)$ which satisfies for every $R_1, R_2 \in (1, \infty)$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ and every random variable $\vartheta: \Omega \rightarrow [-R_2, R_2]^{\mathfrak{d}(\ell)}$ it holds that

$$\begin{aligned} \mathbb{E} \left[\int_D |u(y) - \mathcal{N}_L^{\ell, \vartheta}(y)|^2 dy \right] &\leq c \mathbb{1}_{(R_1, \infty)}(c) + \lambda(D) \mathbb{E} \left[\mathbb{F}_{M, \ell}(\vartheta) - \inf_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} \mathbb{F}_{M, \ell}(\theta) \right] \\ &+ c^L (\mathfrak{d}(\ell))^{4q} \left[\max\{R_1, R_2\} \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\} \right]^{2L} (M_1)^{-1/2} + c \left[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\} \right]^{-\frac{2}{d+5}}. \end{aligned} \quad (157)$$

(cf. Definition 1.1). Note that the fact that for all $x \in (-\infty, c)$ it holds that $\exp(-x) \geq \exp(-c)$ ensures that for all $R \in (0, \infty)$ it holds that

$$\mathbb{1}_{(R, \infty)}(c) = \mathbb{1}_{(-\infty, c)}(R) \leq \exp(c) \exp(-R) \mathbb{1}_{(-\infty, c)}(R). \quad (158)$$

Combining this and (157) implies (156). The proof of Corollary 4.7 is thus complete. \square

Corollary 4.8. *Let $T, \kappa \in (0, \infty)$, $d \in \mathbb{N}$, let $u \in C^2([0, T] \times \mathbb{R}^d, \mathbb{R})$ be at most polynomially growing, assume for all $t \in (0, T)$, $x \in \mathbb{R}^d$ that*

$$\frac{\partial}{\partial t} u(t, x) + \frac{\kappa^2}{2} \Delta_x u(t, x) = 0, \quad (159)$$

let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $\mathbb{T}_n: \Omega \rightarrow [0, T]$, $n \in \mathbb{N}$, be *i.i.d.* random variables, let $\mathbb{X}_n: \Omega \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}$, be *i.i.d.* random variables, let $W^n: [0, T] \times \Omega \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}$, be *i.i.d.* standard Brownian motions, let $D \subseteq (0, T) \times \mathbb{R}^d$ be non-empty, open, and bounded, assume that $(\mathbb{T}_1, \mathbb{X}_1)$ is uniformly distributed on D , assume that $(\mathbb{T}_n)_{n \in \mathbb{N}}$, $(\mathbb{X}_n)_{n \in \mathbb{N}}$, and $(W^n)_{n \in \mathbb{N}}$ are independent, and for every $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ let $\mathbb{F}_{M, \ell}: \mathbb{R}^{\mathfrak{d}(\ell)} \rightarrow \mathbb{R}$ satisfy for all $\theta \in \mathbb{R}^{\mathfrak{d}(\ell)}$ that

$$\mathbb{F}_{M, \ell}(\theta) = \frac{1}{M_1} \sum_{m=1}^{M_1} \left| \mathcal{N}_L^{\ell, \theta}(\mathbb{T}_m, \mathbb{X}_m) - \left[\frac{1}{M_2} \sum_{n=1}^{M_2} u(T, \mathbb{X}_m + \kappa W_{T-\mathbb{T}_m}^{m, M_2+n}) \right] \right|^2 \quad (160)$$

(cf. Definition 1.1). Then there exists $c \in (0, \infty)$ such that for every $R \in (1, \infty)$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ and every random variable $\vartheta: \Omega \rightarrow [-R, R]^{\mathfrak{d}(\ell)}$ it holds that

$$\begin{aligned} \mathbb{E} \left[\int_D |u(y) - \mathcal{N}_L^{\ell, \vartheta}(y)|^2 dy \right] &\leq [R + \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{cL} (M_1)^{-1/2} \\ &+ c \left[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\} \right]^{-2/(d+5)} + \lambda(D) \mathbb{E} \left[\mathbb{F}_{M, \ell}(\vartheta) - \inf_{\theta \in (-c, c)^{\mathfrak{d}(\ell)}} \mathbb{F}_{M, \ell}(\theta) \right] \end{aligned} \quad (161)$$

(cf. Definition 1.1).

Proof of Corollary 4.8. Throughout this proof let $q \in [1, \infty)$ satisfy

$$q = \sup_{x \in \mathbb{R}^d} ((1 + |x|)^{-q} |u(T, x)|). \quad (162)$$

Observe that (162) and Corollary 4.7 show that there exists $c \in (0, \infty)$ which satisfies for every $R_1, R_2 \in (1, \infty)$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ and every random variable $\vartheta: \Omega \rightarrow [-R_2, R_2]^{\mathfrak{d}(\ell)}$ it holds that

$$\begin{aligned} \mathbb{E} \left[\int_D |u(y) - \mathcal{N}_L^{\ell, \vartheta}(y)|^2 dy \right] &\leq c^L (\mathfrak{d}(\ell))^{4q} \left[\max\{R_1, R_2\} \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\} \right]^{2L} (M_1)^{-1/2} + c \exp(-R_1) \mathbb{1}_{(-\infty, c)}(R_1) \\ &+ c \left[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\} \right]^{-\frac{2}{d+5}} + \lambda(D) \mathbb{E} \left[\mathbb{F}_{M, \ell}(\vartheta) - \inf_{\theta \in (-R_1, R_1)^{\mathfrak{d}(\ell)}} \mathbb{F}_{M, \ell}(\theta) \right] \end{aligned} \quad (163)$$

(cf. Definition 1.1). Note that (163) proves that for every $R \in (1, \infty)$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ and every random variable $\vartheta: \Omega \rightarrow [-R, R]^{\mathfrak{d}(\ell)}$ it holds that

$$\begin{aligned} \mathbb{E} \left[\int_D |u(y) - \mathcal{N}_L^{\ell, \vartheta}(y)|^2 dy \right] &\leq c^L (\mathfrak{d}(\ell))^{4q} \left[R \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\} \right]^{2L} (M_1)^{-1/2} \\ &+ c \left[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\} \right]^{-\frac{2}{d+5}} + \lambda(D) \mathbb{E} \left[\mathbb{F}_{M, \ell}(\vartheta) - \inf_{\theta \in (-c, c)^{\mathfrak{d}(\ell)}} \mathbb{F}_{M, \ell}(\theta) \right]. \end{aligned} \quad (164)$$

In the next step we observe that the fact that for all $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ it holds that $\mathfrak{d}(\ell) = \sum_{i=1}^L \ell_i(\ell_{i-1} + 1)$ demonstrates that for all $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ it holds that

$$\begin{aligned} \mathfrak{d}(\ell) &= (d+1)\ell_1 + \left[\sum_{i=2}^{L-1} \ell_i(\ell_{i-1} + 1) \right] + 2\ell_{L-1} \\ &\leq (d+1)[\max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^2 + 2(L-2)[\max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^2 \\ &\quad + 2[\max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^2 \\ &\leq (2L+d)[\max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^2. \end{aligned} \quad (165)$$

This and (164) establish that for every $R \in (1, \infty)$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ and every random variable $\vartheta: \Omega \rightarrow [-R, R]^{\mathfrak{d}(\ell)}$ it holds that

$$\begin{aligned} \mathbb{E}[\int_D |u(y) - \mathcal{N}_L^{\ell, \vartheta}(y)|^2 dy] &\leq c^L (2L+d)^{4q} [R \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{2L+8q} (M_1)^{-1/2} \\ &\quad + c[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{\frac{-2}{d+5}} + \lambda(D) \mathbb{E}[\mathbb{F}_{M, \ell}(\vartheta) - \inf_{\theta \in (-c, c)^{\mathfrak{d}(\ell)}} \mathbb{F}_{M, \ell}(\theta)]. \end{aligned} \quad (166)$$

Hence, we obtain for every $R \in (1, \infty)$, $M = (M_1, M_2) \in \mathbb{N}^2$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ and every random variable $\vartheta: \Omega \rightarrow [-R, R]^{\mathfrak{d}(\ell)}$ that

$$\begin{aligned} \mathbb{E}[\int_D |u(y) - \mathcal{N}_L^{\ell, \vartheta}(y)|^2 dy] &\leq c^L (2L+d)^{4q} [R + \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{4L+16q} (M_1)^{-1/2} \\ &\quad + c[\min\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{\frac{-2}{d+5}} + \lambda(D) \mathbb{E}[\mathbb{F}_{M, \ell}(\vartheta) - \inf_{\theta \in (-c, c)^{\mathfrak{d}(\ell)}} \mathbb{F}_{M, \ell}(\theta)]. \end{aligned} \quad (167)$$

Next, note that the fact that for all $x \in (0, \infty)$ it holds that $2^x \geq x$ ensures that there exist $c_1, c_2 \in (0, \infty)$ such that for all $R \in (1, \infty)$, $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in (\{d+1\} \times \mathbb{N}^{L-1} \times \{1\})$ it holds that

$$\begin{aligned} c^L (2L+d)^{4q} [R + \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{4L+16q} \\ \leq 2^{Lc} 2^{Lc_1} [R + \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{4L+16q} \leq [R + \max\{\ell_1, \ell_2, \dots, \ell_{L-1}\}]^{Lc_2}. \end{aligned} \quad (168)$$

Combining this and (167) implies (161). The proof of Corollary 4.8 is thus complete. \square

4.5. Convergence results for the deep Kolmogorov method

Corollary 4.9. *Let $T, \kappa \in (0, \infty)$, $d \in \mathbb{N}$, let $u \in C^2([0, T] \times \mathbb{R}^d, \mathbb{R})$ be at most polynomially growing, assume for all $t \in (0, T)$, $x \in \mathbb{R}^d$ that*

$$\frac{\partial}{\partial t} u(t, x) + \frac{1}{2} \Delta_x u(t, x) = 0, \quad (169)$$

let $q \in [1, \infty)$ satisfy $\sup_{x \in \mathbb{R}^d} ((1 + \|x\|)^{-q} |u(T, x)|) < \infty$, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $\mathbb{T}_n: \Omega \rightarrow [0, T]$, $n \in \mathbb{N}$, be *i.i.d.* random variables, let $\mathbb{X}_n: \Omega \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}$, be *i.i.d.* random variables, let $W^n: [0, T] \times \Omega \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}$, be *i.i.d.* standard Brownian motions, let $D \subseteq (0, T) \times \mathbb{R}^d$ be non-empty, open, and bounded, assume that $(\mathbb{T}_1, \mathbb{X}_1)$ is uniformly distributed on D , assume that $(\mathbb{T}_n)_{n \in \mathbb{N}}$, $(\mathbb{X}_n)_{n \in \mathbb{N}}$, and $(W^n)_{n \in \mathbb{N}}$ are independent, for every $k \in \mathbb{N}$ let $M^k = (M_1^k, M_2^k) \in \mathbb{N}^2$, $L_k \in \mathbb{N} \setminus \{1\}$, $\ell_k = (\ell_k^0, \ell_k^1, \dots, \ell_k^{L_k}) \in (\{d+1\} \times \mathbb{N}^{L_k-1} \times \{1\})$, $R_k \in [1, \infty)$ and let $\mathbb{F}_k: \mathbb{R}^{\mathfrak{d}(\ell_k)} \rightarrow \mathbb{R}$ satisfy for all $\theta \in \mathbb{R}^{\mathfrak{d}(\ell_k)}$ that

$$\mathbb{F}_k(\theta) = \frac{1}{M_1^k} \sum_{m=1}^{M_1^k} |\mathcal{N}_L^{\ell_k, \theta}(\mathbb{T}_m, \mathbb{X}_m) - [\frac{1}{M_2^k} \sum_{n=1}^{M_2^k} u(T, \mathbb{X}_m + W_{T-\mathbb{T}_m}^{m, M_2^k+n})]|^2, \quad (170)$$

assume $\liminf_{k \rightarrow \infty} \min\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\} = \infty$, assume $\limsup_{k \rightarrow \infty} ((R_k)^{4L_k} [\max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^{4L_k+8q} (M_1^k)^{-1}) = 0$, and let $\vartheta_k: \Omega \rightarrow [-R_k, R_k]^{\mathfrak{d}(\ell_k)}$, $k \in \mathbb{N}$, be random variables which satisfy $\limsup_{k \rightarrow \infty} \mathbb{E}[\mathbb{F}_k(\vartheta_k) - \inf_{\theta \in \mathbb{R}^{\mathfrak{d}(\ell_k)}} \mathbb{F}_k(\theta)] = 0$ (cf. Definition 1.1). Then

$$\limsup_{k \rightarrow \infty} \mathbb{E}[\int_D |u(y) - \mathcal{N}_L^{\ell_k, \vartheta_k}(y)|^2 dy] = 0. \quad (171)$$

Proof of Corollary 4.9. Observe that Corollary 4.7 shows that there exists $c \in (0, \infty)$ which satisfies for all $k \in \mathbb{N}$ that

$$\begin{aligned} \mathbb{E}[\int_D |u(y) - \mathcal{N}_L^{\ell_k, \vartheta_k}(y)|^2 dy] &\leq c^{L_k} (\mathfrak{d}(\ell_k))^{4q} [R_k \max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^{2L_k} (M_1^k)^{-1/2} \\ &+ c[\min\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^{\frac{-2}{d+5}} + \lambda(D) \mathbb{E}[\mathbb{F}_k(\vartheta_k) - \inf_{\theta \in \mathbb{R}^{\mathfrak{d}(\ell_k)}} \mathbb{F}_k(\theta)]. \end{aligned} \quad (172)$$

Note that the fact that for all $L \in \mathbb{N} \setminus \{1\}$, $\ell = (\ell_0, \ell_1, \dots, \ell_L) \in \mathbb{N}^{L+1}$ it holds that $\mathfrak{d}(\ell) = \sum_{i=1}^L \ell_i(\ell_{i-1} + 1)$ proves that for all $k \in \mathbb{N}$ it holds that

$$\begin{aligned} \mathfrak{d}(\ell_k) &= (d+1)\ell_k^1 + \left[\sum_{i=2}^{L_k-1} \ell_k^i (\ell_k^{i-1} + 1) \right] + 2\ell_k^{L_k-1} \\ &\leq (d+1)[\max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^2 + 2(L_k - 2)[\max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^2 \\ &+ 2[\max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^2 \\ &\leq (2L_k + d)[\max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^2. \end{aligned} \quad (173)$$

This and (172) demonstrate for all $k \in \mathbb{N}$ that

$$\begin{aligned} \mathbb{E}[\int_D |u(y) - \mathcal{N}_L^{\ell_k, \vartheta_k}(y)|^2 dy] &\leq c^{L_k} (2L_k + d)^{4q} (R_k)^{2L_k} [\max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^{2L_k+16q} (M_1^k)^{-1/2} \\ &+ c[\min\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^{\frac{-2}{d+5}} + c \mathbb{E}[\mathbb{F}_k(\vartheta_k) - \inf_{\theta \in \mathbb{R}^{\mathfrak{d}(\ell_k)}} \mathbb{F}_k(\theta)] \\ &\leq (2d)^{8q} c^{L_k} (L_k)^{8q} (R_k)^{2L_k} [\max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^{2L_k+8q} (M_1^k)^{-1/2} \\ &+ c[\min\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^{\frac{-2}{d+5}} + \lambda(D) \mathbb{E}[\mathbb{F}_k(\vartheta_k) - \inf_{\theta \in \mathbb{R}^{\mathfrak{d}(\ell_k)}} \mathbb{F}_k(\theta)]. \end{aligned} \quad (174)$$

The fact that $\sup_{n \in \mathbb{N}} 2^n n^{-4q} < \infty$ hence establishes that there exists $c \in (0, \infty)$ such that for all $k \in \mathbb{N}$ with $\max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\} \geq \max\{c, 2\}$ it holds that

$$\begin{aligned} \mathbb{E}[\int_D |u(y) - \mathcal{N}_L^{\ell_k, \vartheta_k}(y)|^2 dy] &\leq c (R_k)^{2L_k} [\max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^{4L_k+8q} (M_1^k)^{-1/2} \\ &+ c[\min\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^{\frac{-2}{d+5}} + \lambda(D) \mathbb{E}[\mathbb{F}_k(\vartheta_k) - \inf_{\theta \in \mathbb{R}^{\mathfrak{d}(\ell_k)}} \mathbb{F}_k(\theta)]. \end{aligned} \quad (175)$$

Combining this, the assumption that

$$\liminf_{k \rightarrow \infty} \min\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\} = \infty, \quad (176)$$

the assumption that $\liminf_{k \rightarrow \infty} \min\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\} = \infty$, and the assumption that $\limsup_{k \rightarrow \infty} \mathbb{E}[\mathbb{F}_k(\vartheta_k) - \inf_{\theta \in \mathbb{R}^{\mathfrak{d}(\ell_k)}} \mathbb{F}_k(\theta)] = 0$ ensures (171). The proof of Corollary 4.9 is thus complete. \square

Corollary 4.10. *Let $T \in (0, \infty)$, $d \in \mathbb{N}$, let $u \in C^2([0, T] \times \mathbb{R}^d, \mathbb{R})$ be at most polynomially growing, assume for all $t \in [0, T]$, $x \in \mathbb{R}^d$ that*

$$\frac{\partial}{\partial t} u(t, x) + \frac{1}{2} \Delta_x u(t, x) = 0, \quad (177)$$

let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $\mathbb{T}_n: \Omega \rightarrow [0, T]$, $n \in \mathbb{N}$, be *i.i.d.* random variables, let $\mathbb{X}_n: \Omega \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}$, be *i.i.d.* random variables, let $W^n: [0, T] \times \Omega \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}$, be *i.i.d.* standard Brownian motions, let $D \subseteq (0, T) \times \mathbb{R}^d$ be non-empty, open, and bounded, assume that $(\mathbb{T}_1, \mathbb{X}_1)$ is uniformly distributed on D , assume that $(\mathbb{T}_n)_{n \in \mathbb{N}}$, $(\mathbb{X}_n)_{n \in \mathbb{N}}$, and $(W^n)_{n \in \mathbb{N}}$ are independent, for every $k \in \mathbb{N}$ let $M^k = (M_1^k, M_2^k) \in \mathbb{N}^2$, $L_k \in \mathbb{N} \setminus \{1\}$, $\ell_k = (\ell_k^0, \ell_k^1, \dots, \ell_k^{L_k}) \in (\{d+1\} \times \mathbb{N}^{L_k-1} \times \{1\})$, $R_k \in (1, \infty)$ and let $\mathbb{F}_k: \mathbb{R}^{\mathfrak{d}(\ell_k)} \rightarrow \mathbb{R}$ satisfy for all $\theta \in \mathbb{R}^{\mathfrak{d}(\ell_k)}$ that

$$\mathbb{F}_k(\theta) = \frac{1}{M_1^k} \sum_{m=1}^{M_1^k} |\mathcal{N}_L^{\ell_k, \theta}(\mathbb{T}_m, \mathbb{X}_m) - [\frac{1}{M_2^k} \sum_{n=1}^{M_2^k} u(T, \mathbb{X}_m + W_{T-\mathbb{T}_m}^{m, M_2^k+n})]|^2, \quad (178)$$

assume $\liminf_{k \rightarrow \infty} \min\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\} = \infty$, assume $\sup_{c \in [1, \infty)} \limsup_{k \rightarrow \infty} ([R_k + \max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^{cL_k} (M_1^k)^{-1}) = 0$, and let $\vartheta_k: \Omega \rightarrow [-R_k, R_k]^{\mathfrak{d}(\ell_k)}$, $k \in \mathbb{N}$, be random variables which satisfy $\limsup_{k \rightarrow \infty} \mathbb{E}[\mathbb{F}_k(\vartheta_k) - \inf_{\theta \in \mathbb{R}^{\mathfrak{d}(\ell_k)}} \mathbb{F}_k(\theta)] = 0$ (cf. Definition 1.1). Then

$$\limsup_{k \rightarrow \infty} \mathbb{E}[\int_D |u(y) - \mathcal{N}_L^{\ell_k, \vartheta_k}(y)|^2 dy] = 0. \quad (179)$$

Proof of Corollary 4.10. Observe that the Cauchy-Schwarz inequality and the fact that for all $k \in \mathbb{N}$ it holds that $R_k > 0$ imply that for all $k \in \mathbb{N}$ it holds that

$$R_k + \max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\} \geq [R_k \max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^{1/2} \quad (180)$$

This and the assumption that $\sup_{c \in [1, \infty)} \limsup_{k \rightarrow \infty} ([R_k + \max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^{cL_k} (M_1^k)^{-1}) = 0$ show that

$$\limsup_{k \rightarrow \infty} ([R_k \max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^{4L_k} (M_1^k)^{-1}) = 0. \quad (181)$$

The assumption that for all $k \in \mathbb{N}$ it holds that $R_k \geq 1$ and the assumption that for all $k \in \mathbb{N}$ it holds that $L_k \geq 1$ therefore prove that

$$\limsup_{k \rightarrow \infty} ((R_k)^{4L_k} [\max\{\ell_k^1, \ell_k^2, \dots, \ell_k^{L_k-1}\}]^{4L_k+8q} (M_1^k)^{-1}) = 0. \quad (182)$$

Combining this and Corollary 4.9 demonstrates (179). The proof of Corollary 4.10 is thus complete. \square

Acknowledgements

This work has been partially funded by the European Union (ERC, MONTECARLO, 101045811). The views and the opinions expressed in this work are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Council (ERC). Neither the European Union nor the granting authority can be held responsible for them. We also gratefully acknowledge the Cluster of Excellence EXC 2044-390685587, Mathematics Münster: Dynamics-Geometry-Structure funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation). Most of the specific formulations in the proofs of this work have been created using [74].

References

- [1] ABDELJAWAD, A., AND GROHS, P. Approximations with deep neural networks in Sobolev time-space. *Anal. Appl. (Singap.)* 20, 3 (2022), 499–541.
- [2] ACKERMANN, J., JENTZEN, A., KRUSE, T., KUCKUCK, B., AND PADGETT, J. L. Deep neural networks with ReLU, leaky ReLU, and softplus activation provably overcome the curse of dimensionality for Kolmogorov partial differential equations with Lipschitz nonlinearities in the L^p -sense. *arXiv:2309.13722* (2023), 52 pages.
- [3] ACKERMANN, J., JENTZEN, A., KUCKUCK, B., AND PADGETT, J. L. Deep neural networks with ReLU, leaky ReLU, and softplus activation provably overcome the curse of dimensionality for space-time solutions of semilinear partial differential equations. *arXiv:2406.10876* (2024), 64 pages.
- [4] ALEJO, M. A., COSSETTI, L., FANELLI, L., MUÑOZ, C., AND VALENZUELA, N. Error bounds for Physics Informed Neural Networks in Nonlinear Schrödinger equations placed on unbounded domains. *arXiv:2409.17938* (2024), 31 pages.
- [5] ANDERSSON, K., ANDERSSON, A., AND OOSTERLEE, C. W. Convergence of a robust deep FBSDE method for stochastic control. *SIAM J. Sci. Comput.* 45, 1 (2023), A226–A255.
- [6] ANDERSSON, K., GNOATTO, A., AND TRILLOS, C. A. G. A deep solver for backward stochastic Volterra integral equations. *arXiv:2505.18297* (2025), 25 pages.

- [7] ARIEL NEUFELD, TUAN ANH NGUYEN, S. W. Deep ReLU neural networks overcome the curse of dimensionality when approximating semilinear partial integro-differential equations. *arXiv:2310.15581* (2023), 36 pages.
- [8] BECK, C., BECKER, S., CHERIDITO, P., JENTZEN, A., AND NEUFELD, A. Deep splitting method for parabolic PDEs. *SIAM J. Sci. Comput.* *43*, 5 (2021), A3135–A3154.
- [9] BECK, C., BECKER, S., GROHS, P., JAAFARI, N., AND JENTZEN, A. Solving the Kolmogorov PDE by means of deep learning. *J. Sci. Comput.* *88*, 3 (2021), Paper No. 73, 28.
- [10] BECK, C., HUTZENTHALER, M., JENTZEN, A., AND KUCKUCK, B. An overview on deep learning-based approximation methods for partial differential equations. *Discrete Contin. Dyn. Syst. Ser. B* *28*, 6 (2023), 3697–3746.
- [11] BECK, C., JENTZEN, A., AND KUCKUCK, B. Full error analysis for the training of deep neural networks. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* *25*, 2 (2022), Paper No. 2150020, 76.
- [12] BENEVENTANO, P., CHERIDITO, P., JENTZEN, A., AND VON WURSTEMBERGER, P. High-dimensional approximation spaces of artificial neural networks and applications to partial differential equations. *arXiv:2012.04326* (2020), 31 pages.
- [13] BERNER, J., DABLANDER, M., AND GROHS, P. Numerically solving parametric families of high-dimensional kolmogorov partial differential equations via deep learning. In *Advances in Neural Information Processing Systems* (2020), H. Larochelle, M. Ranzato, R. Hadsell, M. Balcan, and H. Lin, Eds., vol. 33, Curran Associates, Inc., pp. 16615–16627.
- [14] BERNER, J., GROHS, P., AND JENTZEN, A. Analysis of the generalization error: empirical risk minimization over deep artificial neural networks overcomes the curse of dimensionality in the numerical approximation of Black-Scholes partial differential equations. *SIAM J. Math. Data Sci.* *2*, 3 (2020), 631–657.
- [15] BIAGINI, F., GONON, L., AND WALTER, N. Approximation rates for deep calibration of (rough) stochastic volatility models. *SIAM J. Financial Math.* *15*, 3 (2024), 734–784.
- [16] BICHUCH, M., AND HOU, J. Deep PDE solution to BSDE. *Digit Finance* *6* (2024), 727–758.
- [17] BLECHSCHMIDT, J., AND ERNST, O. G. Three ways to solve partial differential equations with neural network—a review. *GAMM-Mitt.* *44*, 2 (2021), Paper No. e202100006, 29.
- [18] BOKANOWSKI, O., PROST, A., AND WARIN, X. Neural networks for first order HJB equations and application to front propagation with obstacle terms. *Partial Differ. Equ. Appl.* *4*, 5 (2023), Paper No. 45, 36.
- [19] BOUSSANGE, V., BECKER, S., JENTZEN, A., KUCKUCK, B., AND PELLISSIER, L. Deep learning approximations for non-local nonlinear PDEs with Neumann boundary conditions. *Partial Differ. Equ. Appl.* *4*, 6 (2023), Paper No. 51, 51.
- [20] CAI, S., MAO, Z., WANG, Z., YIN, M., AND KARNIADAKIS, G. E. Physics-informed neural networks (PINNs) for fluid mechanics: a review. *Acta Mech. Sin.* *37*, 12 (2021), 1727–1738.
- [21] CARMONA, R., AND LAURIÈRE, M. Convergence analysis of machine learning algorithms for the numerical solution of mean field control and games I: The ergodic case. *SIAM J. Numer. Anal.* *59*, 3 (2021), 1455–1485.

- [22] CASTRO, J. Deep learning schemes for parabolic nonlocal integro-differential equations. *Partial Differ. Equ. Appl.* 3, 6 (2022), Paper No. 77, 35.
- [23] CHAN-WAI-NAM, Q., MIKAEL, J., AND WARIN, X. Machine learning for semi linear PDEs. *J. Sci. Comput.* 79, 3 (2019), 1667–1712.
- [24] CHASSAGNEUX, J.-F., CHEN, J., AND FRIKHA, N. Deep Runge-Kutta schemes for BSDEs. *arXiv:2212.14372* (2022), 33 pages.
- [25] CHEN, F., HUANG, J., WANG, C., AND YANG, H. Friedrichs learning: weak solutions of partial differential equations via deep learning. *SIAM J. Sci. Comput.* 45, 3 (2023), A1271–A1299.
- [26] CHERIDITO, P., AND ROSSMANNEK, F. Efficient Sobolev approximation of linear parabolic PDEs in high dimensions. *arXiv:2306.16811* (2023), 28 pages.
- [27] CUOMO, S., SCHIANO DI COLA, V., GIAMPAOLO, F., ROZZA, G., RAISSI, M., AND PICCIALI, F. Scientific machine learning through physics-informed neural networks: where we are and what’s next. *J. Sci. Comput.* 92, 3 (2022), Paper No. 88, 62.
- [28] DE RYCK, T., AND MISHRA, S. Numerical analysis of physics-informed neural networks and related models in physics-informed machine learning. *Acta Numer.* 33 (2024), 633–713.
- [29] DEREICH, S., GRAEBER, R., AND JENTZEN, A. Non-convergence of Adam and other adaptive stochastic gradient descent optimization methods for non-vanishing learning rates. *arXiv:2407.08100* (2024), 54 pages.
- [30] DO, T., HANNIBAL, S., AND JENTZEN, A. Non-convergence to global minimizers in data driven supervised deep learning: Adam and stochastic gradient descent optimization provably fail to converge to global minimizers in the training of deep neural networks with ReLU activation. *arXiv:2410.10533* (2024), 91 pages.
- [31] DOUMÈCHE, N., BACH, F., BIAU, G., AND BOYER, C. Physics-informed kernel learning. *arXiv:2409.13786* (2024), 35 pages.
- [32] DOUMÈCHE, N., BACH, F., BIAU, G., AND BOYER, C. Physics-informed machine learning as a kernel method. *arXiv:2402.07514* (2024), 52 pages.
- [33] E, W., HAN, J., AND JENTZEN, A. Deep learning-based numerical methods for high-dimensional parabolic partial differential equations and backward stochastic differential equations. *Commun. Math. Stat.* 5, 4 (2017), 349–380.
- [34] E, W., HAN, J., AND JENTZEN, A. Algorithms for solving high dimensional PDEs: from nonlinear Monte Carlo to machine learning. *Nonlinearity* 35, 1 (2022), 278–310.
- [35] E, W., AND YU, B. The deep Ritz method: a deep learning-based numerical algorithm for solving variational problems. *Commun. Math. Stat.* 6, 1 (2018), 1–12.
- [36] EIGEL, M., SCHNEIDER, R., TRUNSCHKE, P., AND WOLF, S. Variational Monte Carlo—bridging concepts of machine learning and high-dimensional partial differential equations. *Adv. Comput. Math.* 45, 5-6 (2019), 2503–2532.
- [37] ELBRÄCHTER, D., GROHS, P., JENTZEN, A., AND SCHWAB, C. DNN expression rate analysis of high-dimensional PDEs: application to option pricing. *Constr. Approx.* 55, 1 (2022), 3–71.

- [38] FEISCHL, M., LASSER, C., LUBICH, C., AND NICK, J. Regularized dynamical parametric approximation. *arXiv:2403.19234* (2024), 38 pages.
- [39] FREY, R., AND KÖCK, V. Convergence analysis of the deep splitting scheme: the case of partial integro-differential equations and the associated forward backward SDEs with jumps. *SIAM J. Sci. Comput.* *47*, 1 (2025), A527–A552.
- [40] GARCÍA-CERVERA, C. J., KESSLER, M., AND PERIAGO, F. Control of partial differential equations via physics-informed neural networks. *J. Optim. Theory Appl.* *196*, 2 (2023), 391–414.
- [41] GERMAIN, M., PHAM, H., AND WARIN, X. Neural networks-based algorithms for stochastic control and PDEs in finance. *arXiv:2101.08068* (2021), 27 pages.
- [42] GERMAIN, M., PHAM, H., AND WARIN, X. Approximation error analysis of some deep backward schemes for nonlinear PDEs. *SIAM J. Sci. Comput.* *44*, 1 (2022), A28–A56.
- [43] GLAU, K., AND WUNDERLICH, L. The deep parametric PDE method and applications to option pricing. *Appl. Math. Comput.* *432* (2022), Paper No. 127355, 21.
- [44] GOBET, E., LÓPEZ-SALAS, J. G., AND VÁZQUEZ, C. Quasi-regression Monte-Carlo scheme for semi-linear PDEs and BSDEs with large scale parallelization on GPUs. *Arch. Comput. Methods Eng.* *27*, 3 (2020), 889–921.
- [45] GONON, L. Random feature neural networks learn Black-Scholes type PDEs without curse of dimensionality. *J. Mach. Learn. Res.* *24* (2023), Paper No. [189], 51.
- [46] GONON, L. Deep neural network expressivity for optimal stopping problems. *Finance Stoch.* *28*, 3 (2024), 865–910.
- [47] GONON, L., GROHS, P., JENTZEN, A., KOFLER, D., AND ŠIŠKA, D. Uniform error estimates for artificial neural network approximations for heat equations. *IMA J. Numer. Anal.* *42*, 3 (2022), 1991–2054.
- [48] GONON, L., JENTZEN, A., KUCKUCK, B., LIANG, S., RIEKERT, A., AND VON WURSTEMBERGER, P. An overview on machine learning methods for partial differential equations: from physics informed neural networks to deep operator learning. *arXiv:2408.13222* (2024), 59 pages.
- [49] GONON, L., AND SCHWAB, C. Deep ReLU network expression rates for option prices in high-dimensional, exponential Lévy models. *Finance Stoch.* *25*, 4 (2021), 615–657.
- [50] GONON, L., AND SCHWAB, C. Deep ReLU neural networks overcome the curse of dimensionality for partial integrodifferential equations. *Anal. Appl. (Singap.)* *21*, 1 (2023), 1–47.
- [51] GRIBONVAL, R., KUTYNIOK, G., NIELSEN, M., AND VOIGTLAENDER, F. Approximation spaces of deep neural networks. *Constr. Approx.* *55*, 1 (2022), 259–367.
- [52] GROHS, P., HORNUNG, F., JENTZEN, A., AND VON WURSTEMBERGER, P. A proof that artificial neural networks overcome the curse of dimensionality in the numerical approximation of Black-Scholes partial differential equations. *Mem. Amer. Math. Soc.* *284*, 1410 (2023), v+93.
- [53] GU, K., FANG, P., SUN, Z., AND DU, R. Error analysis of the mixed residual method for elliptic equations. *Numer. Math-Theory. Me* *17*, 2 (2024), 534–554.

- [54] GÜHRING, I., KUTYNIOK, G., AND PETERSEN, P. Error bounds for approximations with deep ReLU neural networks in $W^{s,p}$ norms. *Anal. Appl. (Singap.)* 18, 5 (2020), 803–859.
- [55] GÜHRING, I., RASLAN, M., AND KUTYNIOK, G. Expressivity of deep neural networks. In *Mathematical aspects of deep learning*. Cambridge Univ. Press, Cambridge, 2023, pp. 149–199.
- [56] GÜHRING, I., AND RASLAN, M. Approximation rates for neural networks with encodable weights in smoothness spaces. *Neural Networks* 134 (2021), 107–130.
- [57] HAIRER, M., HUTZENTHALER, M., AND JENTZEN, A. Loss of regularity for Kolmogorov equations. *arXiv:1209.6035* (2012), 62 pages.
- [58] HAN, J., HU, R., AND LONG, J. Convergence of deep fictitious play for stochastic differential games. *Front. Math. Finance* 1, 2 (2022), 287–319.
- [59] HAN, J., JENTZEN, A., AND E, W. Solving high-dimensional partial differential equations using deep learning. *Proc. Natl. Acad. Sci. USA* 115, 34 (2018), 8505–8510.
- [60] HAN, J., AND LONG, J. Convergence of the deep BSDE method for coupled FBSDEs. *Probab. Uncertain. Quant. Risk* 5 (2020), Paper No. 5, 33.
- [61] HU, R., AND LAURIÈRE, M. Recent developments in machine learning methods for stochastic control and games. *Numer. Algebra Control Optim.* 14, 3 (2024), 435–525.
- [62] HU, Z., SHUKLA, K., KARNIADAKIS, G. E., AND KAWAGUCHI, K. Tackling the curse of dimensionality with physics-informed neural networks. *Neural Networks* 176 (2024), 106369.
- [63] HURÉ, C., PHAM, H., AND WARIN, X. Deep backward schemes for high-dimensional nonlinear PDEs. *Math. Comp.* 89, 324 (2020), 1547–1579.
- [64] HUTZENTHALER, M., JENTZEN, A., KRUSE, T., AND NGUYEN, T. A. A proof that rectified deep neural networks overcome the curse of dimensionality in the numerical approximation of semilinear heat equations. *Partial Differ. Equ. Appl.* 1, 2 (2020), Paper No. 10, 34.
- [65] HUTZENTHALER, M., JENTZEN, A., KUCKUCK, B., AND PADGETT, J. L. Strong L^p -error analysis of nonlinear Monte Carlo approximations for high-dimensional semilinear partial differential equations. *arXiv:2110.08297* (2021), 42 pages.
- [66] HUTZENTHALER, M., JENTZEN, A., POHL, K., RIEKERT, A., AND SCARPA, L. Convergence proof for stochastic gradient descent in the training of deep neural networks with ReLU activation for constant target functions. *arXiv:2112.07369* (2021), 71 pages.
- [67] JENTZEN, A., KUCKUCK, B., AND VON WURSTEMBERGER, P. Mathematical Introduction to Deep Learning: Methods, Implementations, and Theory. *arXiv:2310.20360* (2025), 737 pages.
- [68] JIANG, Y., AND LI, J. Convergence of the deep BSDE method for FBSDEs with non-Lipschitz coefficients. *Probab. Uncertain. Quant. Risk* 6, 4 (2021), 391–408.
- [69] JIAO, Y., LAI, Y., LO, Y., WANG, Y., AND YANG, Y. Error Analysis of Deep Ritz Methods for Elliptic Equations. *arXiv:2107.14478* (2021), 23 pages.
- [70] JIAO, Y., LAI, Y., WANG, Y., YANG, H., AND YANG, Y. Convergence Analysis of the Deep Galerkin Method for Weak Solutions. *arXiv:2302.02405* (2023), 21 pages.

- [71] JIAO, Z., JIA, W., AND ZHU, W. Solving a class of stochastic optimal control problems by physics-informed neural networks. *arXiv: 2402.15592* (2024), 8 pages.
- [72] JOVANOVIĆ, B. S. S., AND SÜLI, E. *Analysis of finite difference schemes*, vol. 46 of *Springer Series in Computational Mathematics*. Springer, London, 2014. For linear partial differential equations with generalized solutions.
- [73] KNOCHENHAUER, C., HAGER, O., REIMERS, C., SCHNELL, L., AND WÜRSCHMIDT, M. Convergence Rates for a Deep Learning Algorithm for Semilinear PDEs. *SSRN Electronic Journal* (2021).
- [74] KUCKUCK, B. Some useful latex commands. <https://latex.bennokuckuck.de> [Accessed 22-August-2025] (2025).
- [75] KUTYNIOK, G., PETERSEN, P., RASLAN, M., AND SCHNEIDER, R. A theoretical analysis of deep neural networks and parametric PDEs. *Constr. Approx.* 55, 1 (2022), 73–125.
- [76] LU, J., LU, Y., AND WANG, M. A Priori Generalization Analysis of the Deep Ritz Method for Solving High Dimensional Elliptic Partial Differential Equations. *Proc. Mach. Learn. Res.* 134 (2021), 1–46.
- [77] LU, J., SHEN, Z., YANG, H., AND ZHANG, S. Deep network approximation for smooth functions. *SIAM J. Math. Anal.* 53, 5 (2021), 5465–5506.
- [78] LUO, T., AND YANG, H. Two-Layer Neural Networks for Partial Differential Equations: Optimization and Generalization Theory. *arXiv:2006.15733* (2020), 31 pages.
- [79] MAO, T., AND ZHOU, D.-X. Rates of approximation by ReLU shallow neural networks. *J. Complexity* 79 (2023), Paper No. 101784, 21.
- [80] MISHRA, S., AND MOLINARO, R. Estimates on the generalization error of physics-informed neural networks for approximating PDEs. *IMA J. Numer. Anal.* 43, 1 (2023), 1–43.
- [81] MUÑOZ, C., AND VALENZUELA, N. Bounds on the approximation error for deep neural networks applied to dispersive models: Nonlinear waves. *arXiv:2405.13566* (2024), 50 pages.
- [82] NEUFELD, A., AND NGUYEN, T. A. Rectified deep neural networks overcome the curse of dimensionality in the numerical approximation of gradient-dependent semilinear heat equations. *arXiv:2403.09200* (2024), 25 pages.
- [83] NEUFELD, A., AND NGUYEN, T. A. Rectified deep neural networks overcome the curse of dimensionality when approximating solutions of McKean-Vlasov stochastic differential equations. *J. Math. Anal. Appl.* 541, 1 (2025), Paper No. 128661, 25.
- [84] NEUFELD, A., AND SCHMOCKER, P. Universal approximation property of Banach space-valued random feature models including random neural networks. *arXiv:2312.08410* (2023), 44 pages.
- [85] NEUFELD, A., SCHMOCKER, P., AND WU, S. Full error analysis of the random deep splitting method for nonlinear parabolic PDEs and PIDEs. *arXiv:2405.05192* (2024), 42 pages.
- [86] OPSCHOOR, J. A. A., PETERSEN, P. C., AND SCHWAB, C. Deep ReLU networks and high-order finite element methods. *Anal. Appl. (Singap.)* 18, 5 (2020), 715–770.

- [87] OUYANG, D., XIAO, J., AND WANG, X. Generalization Error Analysis of Deep Backward Dynamic Programming for Solving Nonlinear PDEs. [arXiv:2407.14566](#) (2024), 29 pages.
- [88] PENG, Y., WEI, P., AND WEI, W. Deep Penalty Methods: A Class of Deep Learning Algorithms for Solving High Dimensional Optimal Stopping Problems. [arXiv:2405.11392](#) (2024), 27 pages.
- [89] PETERSEN, P., AND VOIGTLAENDER, F. Optimal approximation of piecewise smooth functions using deep ReLU neural networks. *Neural Networks* 108 (2018), 296–330.
- [90] RAISSI, M., PERDIKARIS, P., AND KARNIADAKIS, G. E. Physics-informed neural networks: a deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. *J. Comput. Phys.* 378 (2019), 686–707.
- [91] REISINGER, C., AND ZHANG, Y. Rectified deep neural networks overcome the curse of dimensionality for nonsmooth value functions in zero-sum games of nonlinear stiff systems. *Anal. Appl. (Singap.)* 18, 6 (2020), 951–999.
- [92] SCHMIDT-HIEBER, J. Nonparametric regression using deep neural networks with ReLU activation function. *Ann. Statist.* 48, 4 (2020), 1875–1897.
- [93] SCHMIDT-HIEBER, J. Supplement to "Nonparametric regression using deep neural networks with ReLU activation function". [arXiv:1708.06633](#) (2020), 16 pages.
- [94] SIRIGNANO, J., AND SPILIOPOULOS, K. DGM: a deep learning algorithm for solving partial differential equations. *J. Comput. Phys.* 375 (2018), 1339–1364.
- [95] STEIN, E. M. *Singular integrals and differentiability properties of functions*, vol. No. 30 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1970.
- [96] THOMÉE, V. *Galerkin finite element methods for parabolic problems*, second ed., vol. 25 of *Springer Series in Computational Mathematics*. Springer-Verlag, Berlin, 2006.
- [97] VALENZUELA, N. A new approach for the fractional Laplacian via deep neural networks. [arXiv:2205.05229](#) (2022), 55 pages.
- [98] WANG, W., WANG, J., LI, J., GAO, F., AND FU, Y. Deep learning numerical methods for high-dimensional fully nonlinear PIDEs and coupled FBSDEs with jumps. [arXiv:2301.12895](#) (2023), 26 pages.
- [99] WANG, Y., NI, Y.-H., CHEN, Z., AND ZHANG, J.-F. Probabilistic framework of Howard’s policy iteration: BML evaluation and robust convergence analysis. *IEEE Trans. Automat. Control* 69, 8 (2024), 5200–5215.
- [100] WANG, Z., AND TANG, S. Gradient convergence of deep learning-based numerical methods for BSDEs. *Chinese Ann. Math. Ser. B* 42, 2 (2021), 199–216.
- [101] XIAO, J., FU, F., AND WANG, X. Deep Learning Based on Randomized Quasi-Monte Carlo Method for Solving Linear Kolmogorov Partial Differential Equation. [arXiv:2310.18100](#) (2023), 38 pages.
- [102] XIAO, J., AND WANG, X. Error analysis for empirical risk minimization over clipped ReLU networks in solving linear Kolmogorov partial differential equations. [arXiv:2310.12582](#) (2023), 28 pages.
- [103] YAROTSKY, D. Error bounds for approximations with deep ReLU networks. *Neural Networks* 94 (2017), 103–114.

- [104] YE, Z., AND WANG, W. FBSJNN: A Theoretically Interpretable and Efficiently Deep Learning method for Solving Partial Integro-Differential Equations. [arXiv:2412.11010](https://arxiv.org/abs/2412.11010) (2024), 29 pages.